### 2016 FISHING FORECAST AND TIPS



Welcome to the 2016 Fishing Forecast for Kentucky's major fisheries. The forecast is based on 2015 fish population surveys, creel surveys, fish stockings, and historical knowledge of the fisheries. This handout is designed to assist anglers in planning their fishing trips and improving their fishing success. Additional fishing information is available from the Department's website at <a href="fw.ky.gov">fw.ky.gov</a> or by obtaining copies of the 2016 Sport Fishing and Boating Guide available at most sporting goods stores. The Kentucky Trout Waters brochure is contained in the 2016 Fishing and Boating Guide.

To locate fishing access sites in Kentucky, visit our website and click on "Fish" and then "Where to Fish". You will be able to search for your favorite water bodies and get directions to all major boat launches and access sites.

Opening day of the 2016 fishing season starts March 1 with the new year's license, so take a trip to your local sporting good store, get online at fw.ky.gov, or call 1-877-598-2401 to purchase your 2016 fishing license.

The Fishing Forecast was partially financed through funds provided by your purchase of fishing equipment and motor boat fuels under the Federal Sport Fish Restoration Program.





#### **New and Expanding Fisheries in 2016**

- Saugeye are beginning to reach harvestable size (>15 inches) at Guist Creek, Bullock Pen and A.J.
   Jolly lakes. Potential harvestable fish at Boltz Lake in fall of 2016.
- The new lower section of Hatchery Creek at the Wolf Creek National Fish Hatchery is scheduled to open late spring 2016. Check the KDFWR website (fw.ky.gov) for opening date updates.
- An ongoing major fish habitat project at Cave Run Lake should provide increased fishing
  opportunities in the areas of the newly established habitat. See the KDFWR website (fw.ky.gov) for
  GPS locations of the new habitat structures.
- Improved fishing access has been constructed at Meldahl Lock and Dam. Improvements include a fishing pier, grouted walkway, parking and restrooms.

#### **Fishing Forecast Cheat Sheet**

Make sure you check out the new Fishing Forecast Cheat Sheet located at the end of this publication. For those looking for a quick answer to where the best fishing can be found for each species, the Cheat Sheet is your guide. The Cheat Sheet lets you know which lakes are forecasted to provide good to excellent fishing for each of your favorite fish species. Just look up the species you are interested in and then see which lakes have a check mark next to them. These are the lakes which will provide the best chance for a memorable fishing trip for that species. The Cheat Sheet is located on two pages, so if you don't see the species you are interested in on the first page; make sure to check the following page.

#### **New Up-and-Comers**

Several lakes and rivers have shown improvements in their fisheries in the last year. You might try and get in on the action at the following water bodies:

- Largemouth bass at Carr Creek Lake (Knott County), Dewey Lake (Floyd County) and McNeely Lake (Jefferson County).
- Bluegill at Briggs Lake (Logan County), Pennyrile Lake (Christian County), Spurlington Lake (Taylor County) and West Fork Drakes Creek Lake (Simpson County).
- Redear sunfish at Briggs Lake (Logan County).
- Crappie at Carr Creek Lake (Knott County) and Taylorsville Lake (Anderson, Nelson and Spencer counties).
- Striped bass at Kentucky Lake Tailwater (Livingston County).
- Hybrid striped bass in Barren River (from Barren River Lake Dam to Green River confluence)
- Blue catfish at Fishtrap Lake (Pike County).

#### Late Winter/Early Spring Fishing Frenzy

As warm winds begin to flow over Kentucky, put your new fishing license to good use by trying a few early spring fishing hot spots. In March, fish activity begins in the smaller bodies of water such as farm ponds and smaller public lakes as water temperatures reach the 50's. These smaller, shallower bodies of water warm quicker than reservoirs in spring. Late winter and early spring may be the best time of the year to catch big fish.

Trophy largemouth bass, especially females, are at their heaviest weight of the entire year during late winter and early spring. Female bass feed heavily during this time to provide fuel for the upcoming spawn. Farm ponds and public lakes are prime spots for early spring largemouth bass fishing before the larger bodies of water warm enough for bass to feed.

A few of the best lakes for catching trophy bass in spring are Lake Barkley, Kentucky Lake and Lake Malone in western Kentucky along with Cedar Creek, Kincaid and Guist Creek lakes in central Kentucky. Greenbo Lake in northeast Kentucky and Wood Creek Lake in southeast Kentucky both produced multiple state record largemouth bass in the past. Woods Creek Lake is the home of the current 13-pound, 10.4 ounce state record. Barren River Lake in west-central Kentucky, Lake Beshear and Lake Mauzy in west Kentucky along with Lake Herrington and Green River lakes in central Kentucky also offer excellent early spring bass fishing. Dewey Lake, Fishtrap and Fishpond lakes give east Kentucky anglers a great shot for an early spring trophy largemouth bass.

If you are after smallmouth bass, head to Green River, Kentucky, Cumberland, Laurel River, Dale Hollow and Fishtrap lakes for a good chance at catching the smallmouth bass of a lifetime in late winter and early spring. Streams also provide excellent fishing for smallmouth bass in early spring throughout central and eastern Kentucky. Adult fish migrate from their winter habitat in deep, slow moving holes in the lower sections of streams to spawning areas in smaller tributary streams. Elkhorn Creek, South Fork, Licking River and several upper Kentucky, Green and Barren River tributaries offer the best spring fishing opportunities for smallmouth bass.

Slab crappie are ready to be caught, beginning in late winter, at many reservoirs in central and western Kentucky. Catches of crappie over 12 inches are not uncommon at Barkley and Kentucky lakes during this time. Black crappie move into shallow water well before the spawning season in early March, so start early for these fish. Other good bets for crappie include Herrington, Taylorsville, Cumberland, Dewey, Rough, Nolin, Barren and Buckhorn lakes as well as Ohio River embayments.

Excellent white bass fishing can be found during the early spring in the upper sections of reservoirs such as Barkley, Cave Run, Nolin River and Herrington lakes. The Kentucky, Green and Ohio rivers offer excellent spring white bass fishing.

During late winter, sauger and walleye fishing peaks in tailwaters below dams on the Ohio, lower Kentucky, Cumberland and Tennessee rivers. Lake Cumberland, Laurel River and Carr Creek lakes possess excellent walleye fisheries.

If you are interested in muskellunge, they move in spring to shallow shoreline structure and headwater areas in search of warmer water at Cave Run, Green, and Buckhorn lakes. Remember, a 36-inch minimum size limit and one fish daily creel limit is in effect for muskellunge at these three reservoirs.

Do not forget to take advantage of trout waters such as seasonal catch-and-release trout streams, lakes stocked with trout in winter months, Fishing in Neighborhood (FINs) lakes stocked with trout and the Cumberland River tailwater below Lake Cumberland in late winter and early spring. Trout bite willingly in cold water. Expect fishing to improve on the Cumberland Tailwater now that the construction on Wolf Creek Dam is completed.

Try these late winter and early spring hot spots. Not only will they cure your cabin fever, but you might catch the fish of your lifetime.

### **2016 FISHING FORECAST**

Kentucky Department of Fish and Wildlife Resources

The fishing forecast is based on 2015 fish population surveys, planned fish stockings, and projections for 2016.

Lo	ocation/Species	Forecast	Comments
	ARKLEY LAKE and 1		5,600 acres (Trigg, Lyon and Livingston counties)
<u> </u>	Asian carp (silver, bid excellent bowfishing are among the larger rough species often "trash fish" and preparties and dispersion of the ranglers and dispersion of the ranglers and dispersion of the rangle of the	ghead, and grass opportunity and a st fish you will find congregate in the aration instruction spose of fish carc	s carp) are present in the lake and tailwaters. Although unwanted, this has created an a way that anglers can help remove these unwanted fish from our waters. Bighead carp d in north America. The best fishing will be in the tailwaters where Asian carp and other espring and summer. Asian carp make excellent table fare despite their reputation as a his can be found on the KDFWR webpage (fw.ky.gov). Remember to be courteous to easses properly. Harvest of Asian carp is encouraged!!
	Blue catfish	Good/Excellent	Good numbers of 10-20 pounders with the occasional 50+ pounder. Best fishing is during late spring and summer along deep river channels. Look for good flow in the lake for best fishing conditions. Blue catfish can be caught year-round in the tailwater. Fresh caught shad or skipjack is best, but cut-up pieces of Silver carp also works well.
	Bluegill	Good/Excellent	Good numbers of fish up to 9 inches; the best fishing comes in May on sand or gravel spawning beds. Crickets and worms fished under a cork during the spawn is a great option. Fish move deeper as the water warms in summer.
	Channel catfish	Excellent	Best fishing is during May along rocky banks and during summer along the main river channel and adjacent flats. Try stink-bait in shallow water in the spring. Channel catfish can also be caught year-round in the tailwater.
	Crappie	Excellent	Barkley Lake is a little tougher fishing for crappie than Kentucky Lake, though the numbers of crappie are good. You can expect to catch a lot of short fish this year. Many fishermen are changing tactics, casting or trolling light jigs and roadrunners year-round.
	Largemouth bass	Excellent	Shallow water fishing for spawning bass can be amazing in the spring around button ball bushes with floating baits, buzzbaits, spinnerbaits, and plastic creature baits. During post spawn most bass move to drop-offs in the main lake and mouths of the bays. Try fishing Carolina rigs, jigs, and crankbaits in deeper water when there is good current moving through the reservoir. During summertime, fishing shallow near structure can also be effective, particularly at dawn or dusk. In fall and winter, bass move into the bays to feed on shad.
	Redear sunfish	Excellent	Good numbers of 8-10 inch fish with trophies up to 12 inches. Prime fishing occurs in late April and throughout May as fish are spawning. Fish can be found in areas with gravel substrate and vegetation. Redear generally spawn a bit deeper than bluegill.
	Sauger	Poor	Population consists of low numbers of harvestable-size (>14 inch) fish. Many fish are caught trolling deep diving crankbaits near secondary channel breaks during the early summer. Sauger can also be caught while fishing for catfish along the main river channel. Fair fishing is found in the tailwaters.
	Smallmouth bass	Fair	Smallmouth bass make up a small portion of the total bass population but are common in certain habitats. Fishing is best around rocky shorelines near deep water, main lake points, and the Canal during late winter and early spring with suspending jerk baits.
	Striped bass	Good	The tailwater fishing is best for consistent catches, but the lake produces good catches as well. Fish deep main channel ledges near bridges and other places with current. For the best luck try trolling crankbaits or fishing vertically with large jigs. Good trophy potential.

Location/Species	Forecast	Comments
<b>BARKLEY LAKE and</b>	TAILWATER (coi	ntinued)
White bass	Good/Excellent	Fishing success has been decent the last few years. Look for schools of shad breaking the surface during summer and fall. If you can't find them working the surface, fish deeper along ledges on the main lake with a spoon or inline spinner. Trolling the ledges with lighter colored crankbaits can be a lot of fun, especially for children and inexperienced anglers. Similar to striped bass, good fishing occurs in the tailwaters.

		Region Diver Lake tollwater area bybrid fishing is beet, especially during at following
Hybrid striped bass	Good/Excellent	Barren River Lake tailwater area hybrid fishing is best, especially during or following periods of heavy lake discharges. Good fishing also below instream dams at Bowling Green and Greencastle.
Muskellunge	Good	Target mouth of feeder creeks and around wood structure or other current breaks fro dam to Green River confluence. Stocked annually.
Smallmouth bass	Good/Excellent	Good numbers of 12-inch fish. Trophy fish (20-inch plus) available. Fishing best frodam to mouth of Drakes Creek. Target areas above and below riffle/shoal areas and rocky outcroppings.
Spotted bass	Fair	Found throughout Barren River. Fish slower moving pools with rocky banks and logs Fish spinnerbaits and crankbaits in the spring and early summer. Use buzzbaits dur summer.
Walleye/Sauger	Fair/Good	Fishing best below instream dams (Barren River Lake, Greencastle and Bowling Greespecially during December – March. Good-sized walleye (15-30 inch) available. Sauger were stocked from 2010-2014.
RREN RIVER (abov	ve lake upstream	from Long Creek confluence)
Bass	Fair	Largemouth, spotted and smallmouth all found here in fair numbers and good sizes. Largemouth best below 100 bridge; smallmouth best above Coruth or Dunns ford/Celsor Road.
Channel catfish	Good	Best action during early onset of rising water conditions anytime of year, but especial during summer months.

Fair/Good

tailed grubs.

Walleye

Males hanging near riffle areas starting mid-February until early April. Fish deepest section of pools rest of the year with bottom bumping spinners, blade baits or curly

Location/Species	Forecast	Comments
BARREN RIVER LAKE	10,000 acres	(Allen and Barren counties)
		rember) oxygen levels are too low to support fish below 15-18 feet.
Bluegill (sunfish)	Good	Good numbers of 6–8 inch fish. Fish stump beds and back of coves during the spring and fall. Fish deeper in the summer near cove mouths and along bluff walls near laydowns or standing timber.
Catfish	Excellent	Good populations of channel and flathead catfish present with good numbers of larger individuals. Fishing best near creek channels and at night. Blue catfish have been stocked annually since 2010, with some of these already over 30 inches. Fish for blues with live or cut bait in open water areas with jugs or rod and reel.
Crappie	Good/Excellent	Excellent numbers of crappie (black and white) available. Good numbers of larger white crappie (10-inch plus), but few larger black crappie (10-inch plus) available. Ratio of black to white crappie is about 50-50. Target shallower water to find black crappie in spring and early summer. Summer and winter fishing is best around deeper structure in creek/river channel and fish attractor/brush sites.
Hybrid striped bass	Good	Overall numbers are good with large fish (20-inch plus) available. Spring fishing best in upper lake portions at 1st and 2nd riffle areas with spinners, crank baits and curly tailed jigs. Target main lake (around islands in mid-lower lake) early in the morning and at dusk in the summer and fall as schools appear throughout the lake with crankbaits and spoons. 200,000 stocked annually since 2000. Higher numbers of 20 inch-plus available from elevated stocking of 400,000 in 2013.
Largemouth bass	Excellent	Good numbers overall and good numbers of 15-inch plus and 20-inch plus fish.
Smallmouth bass	Fair	Low density population overall; however, quality/trophy fish (20-inch plus) available. Target lower 1/3 of the lake near the dam, fishing rock outcroppings and rocky transition areas.
Spotted bass	Fair	Lower density population overall; however, quality fish (15-inch plus) available. Bridge pylons and bluff walls with rock slides are good places to target spots.
White bass	Fair	Numbers remain low. Best opportunity is during spawning run in late March to mid April with in-line spinners, small jigs and small crankbaits in Barren River, Peter, Skaggs and Beaver creeks above the lake.
Yellow bass	Excellent	Abundant and willing to bite. Most fish are 7-9 inches in length. Bite readily on small jigs, spinners and minnows. Bridge pylons excellent areas to target for summertime bite.

BEAVER LAKE	158 acres	(Anderson County)
Bluegill	Good	High numbers of 6-8 inch fish; ≥ 8 inch fish present.
Channel catfish	Good	Catchable-size channel catfish stocked last year; fish over 15 inches common. Fall fishing good in shallows in coves.
Crappie	Fair/Good	Small population; good numbers of fish between 8-10 inches; larger fish possible.
Largemouth bass	s Good	Good numbers of bass up to 15 inches.
Redear sunfish	Fair	7-9 inch fish available around vegetation; good numbers of 10-11 inch fish possible. Overall, numbers are down due to the significant reduction of aquatic vegetation in lake. 35,625 redear sunfish were stocked in fall of 2015.

Location/Species	Forecast	Comments
BENJY KINMAN LAKI	E 88 acres	(Henry County)
Bluegill	Fair	Fair numbers of 5-7 inch fish.
Channel catfish	Fair	Channel catfish present.
Crappie	Fair/Good	Good numbers of fish between 8-10-inches; larger fish possible.
Largemouth bass	Fair/Good	Good numbers of bass present; very good potential for quality size fish and bass over 20 inches. Trophy-sized ( $\geq$ 23 inch) are present. <b>Note: largemouth bass are catch and release only.</b>
Redear sunfish	Fair	Low numbers of 6-8 inch fish; key in on few areas where aquatic vegetation is present
BERT COMBS LAKE	36 acres	(Clay County)
		uring part of the year due to repairs to the dam.
Bluegill	Good	Good population of bluegill with some large ones mixed in.
Channel catfish	Fair	Stocked every other year.
Largemouth bass	Good	Good numbers of 11-13 inchers with a few large fish.
Rainbow trout	Good	Stocked annually. Fishing pier provides good access for bank anglers targeting trout.
	•	dwell and Christian counties)
Blue catfish	Fair	Blue catfish are being stocked every third year. Their population is doing fair, with good growth. Some already measure over 25 inches and weigh over 5 pounds.
Bluegill	Fair	Numerous small fish with occasional 8-inch fish caught. Boat fishing is the most popul method, but some bank fishing opportunities are also available.
Channel catfish	Excellent	High numbers of 10-15 inch channel catfish in the population. Channel catfish are stocked every other year at 10-12 inches. This is an underutilized fishery. Most angle fishing for catfish use jug lines or noodles. Catfish are one of the reasons people go to Lake Beshear to fish.
Crappie	Fair	Numerous fish in the 6-8 inch range, with an occasional 10 incher. Best fishing during spring in areas of aquatic vegetation and around woody structure. The population is considered stunted; therefore it is ok to harvest the small crappie.
Largemouth bass	Excellent	One of the main reasons anglers fish Lake Beshear. The bass population is dominated by 12-18 inch fish. Best fishing during spring when fish move to shallow water, along the water willow and lily pads. Try fishing the edges of these weeds with a pearl color fluke using a short jerk retrieve, or use a white spinnerbait. Worms, creature baits, an other soft plastics can also be effective during the spawn. In late afternoon, or on cloud days, try top water baits. During summertime, as temperatures heat up, bass often suspend near deep water ledges. These fish can be caught on dropshot rigs or long lined crankbaits and jerkbaits.

Location/Species	Forecast	Comments
BEULAH LAKE 8	7 acres (Jac	ckson County)
Bluegill	Fair	Moderate numbers of 6-7 inch bluegill.
Channel catfish	Fair	Stocked every other year.
Largemouth bass	Good	Abundant 10-13-inch fish with a few large fish (18-inch plus). Good lake to introduce a kid to bass fishing, since bass are fairly abundant.
Rainbow trout	Good	Stocked annually.
Smallmouth bass	Poor	Starting to see a few in the lake.
BOLTZ LAKE 92 a	cres (Grant C	Sounty)
Bluegill	Fair/Good	Good numbers of fish between 6-8 inches.
Catfish	Fair/Good	Catchable-size channel catfish stocked in summer of 2015. Good numbers of channel catfish over the 12 inch size limit. Low numbers of blue catfish; last stocking of catchable-size blue catfish in 2013; 25-inch plus blue catfish possible.
Crappie	Fair/Good	Abundant; majority of fish between 6-8 inches; larger fish possible.
Largemouth bass	Good	Good numbers of bass at and above 12-inch size limit; some potential for bass ≥20 inches.
Redear sunfish	Poor	Low numbers of 6-8 inch fish available around vegetation. 20,775 redear sunfish were stocked in fall of 2015.

#### **BRIGGS LAKE**

Fair

Saugeye

RIGGS LAKE 18 acres (Logan County)
Summertime (mid-June through September) oxygen levels are too low to support fish below 7 feet.

2016.

Channel catfish Good All sizes available. Stocked every other year.  Crappie Fair Large fish present; fish laydown trees, stakebeds and brushpiles in late winter and early spring.  Largemouth bass Fair Better than average numbers of 12-inch plus fish.	Bluegill	Good/Excellent	Good numbers of 8-inch fish available.
spring.	Channel catfish	Good	All sizes available. Stocked every other year.
Largemouth bass Fair Better than average numbers of 12-inch plus fish.	Crappie	Fair	Large fish present; fish laydown trees, stakebeds and brushpiles in late winter and early spring.
	Largemouth bass	Fair	Better than average numbers of 12-inch plus fish.
Redear sunfish Good/Excellent Good numbers of larger fish (8-inch plus) available. Larger fish very shallow in early spring; then move to thermocline depth.	Redear sunfish	Good/Excellent	
Warmouth Good Good numbers of 7-8 inch fish. Fish in/near vegetation.	Warmouth	Good	Good numbers of 7-8 inch fish. Fish in/near vegetation.

First year of stocking; possible harvestable-size saugeye may be present by fall of

Location/Species	Forecast	Comments
<b>BUCKHORN LAKE</b> ar	nd TAILWATER	1,230 acres (Leslie and Perry counties)
Bluegill	Good	Good size distribution and very good numbers; larger fish 7-8 inches. Upper lake areas and heads of coves hold best numbers during summer pool. Shallow mudflats in main lake areas can hold large congregations of fish on spawning beds.
Catfish	Excellent	High numbers of channel catfish with good recruitment of young fish each year. Some flathead catfish available up to 40-50 pounds. Tailwater provides additional good fishery.
Crappie	Good	July and August can provide some very good fishing over shallow mud flats adjacent to main lake channel. Try trolling bait tipped jigs or small crankbaits over the flats. Also, October-November during drawdown to winter pool can be very good in the same locations. Most of legal fish are 9-12 inches with occasional fish to 14 inches.
Largemouth bass	Fair	Excellent growth rates of young bass and fair numbers of 15-17 inch fish with good numbers for fish greater than 20 inches. Spring fishing success can be very good if high water events do not occur.
Muskellunge	Fair	Overall numbers are good, but anglers still contending with majority being smaller fish. Fishing can be difficult for periods following flood events. From January-March fish will congregate in the lower lake and near dam. Trace Branch area has good fishing in July and August. September will find many large fish on edges of weed beds in backs of coves in shallow water. Bank fishing opportunities are good at the tailwater in March and October.
White bass	Poor	Fish range primarily from 8-15 inches and numbers remain low.
BULLOCK PEN LAKE	134 acres	(Grant County)
Bluegill	Poor	Most fish are 6 inches or less, but are abundant.
Catfish	Fair/Good	Catchable-size channel catfish stocked last year; good numbers over the 12-inch size limit and potential for larger fish (≥20 inches). Catchable-size blue catfish stocked each year; 25-inch plus fish possible. Flathead catfish present.
Crappie	Fair	Abundant population of small-size fish (mostly 6-7 inches); few larger fish are present.
Largemouth bass	Good/Excellent	Good numbers of bass at and above size limit (12-inch size limit). Excellent potential for quality-size fish and bass over 20 inches.
Redear sunfish	Poor	Very low numbers.
Saugeye	Fair	Three years of stocking have been completed; harvestable-sized saugeye are present.
White bass	Poor/Fair	Small population of large fish in the 14-17 inch range.
CANNON CREEK LAI	KE 243 acres	(Bell County)
Channel catfish	Fair	Stocked every other year.
Largemouth bass	Fair	Sparse population of mostly 11-13-inch bass in this clear lake.
Rainbow trout	Good	Stocked annually. Good cold water habitat in this deep, clear lake.
Smallmouth bass	Poor	Low density population.
Spotted bass	Fair	Fair population of 10-13-inch fish.

Location/Species	Forecast	Comments
CARPENTER LAKE	68 acres	(Daviess County)
Bluegill	Fair/Good	Good number of 6-7 inch bluegill, but few over 8 inches.
Channel catfish	Good	Stocked every other year. Excellent growth rates produce good numbers of 15-20 inch and larger fish.
Largemouth bass	Fair/Good	Several smaller (8-11 inch) fish in the population, but the number of 12-20 inch fish is decent and holding steady.
Redear sunfish	Fair	Low numbers, but improving. Most 7-8 inches with a few 9-11 inches.

#### CARR CREEK LAKE 710 acres (Knott County)

Hydrilla was documented in Carr Creek Lake for the first time in 2008. Boaters are asked to take measures to prevent the spread of this damaging non-native invasive species within the lake and between lakes by properly inspecting and disinfecting boats, trailers and equipment.

Bluegill	Fair	Good numbers of small bluegill present. Shallow water areas in creek arms provide good fishing in spring and early summer. Larger fish will hold around weedbed edges in summer. Fishing pier near ramp at marina holds fish in submerged brush when lake is at summer pool.
Crappie	Good	Larger black crappie from 9-12 inches and white crappie from 10-16 inches. Deadfall trees and submerged brush piles will congregate large numbers of fish during most of the year.
Largemouth bass	Good	Good numbers of large fish over 20 inches. March-early April anglers have very good fishing success for big fish. Poor recruitment of small fish usually leads to Department stocking of fingerlings each year.
Smallmouth bass	Poor	Low recruitment; a few fish sampled up to 20 inches. Lower half of lake holds best numbers.
Spotted bass	Good	Best numbers are concentrated in lower and mid lake; a few large fish up to 18 inches; excellent numbers through 16 inches.
Walleye	Excellent	Very high numbers of fish from 15-26 inches; a few fish up to 28 inches. March through May are good months to try during day or night. Standing timber will hold suspended fish in summer as well as deeper open water areas with schools of alewife that walleye will suspend beneath.

#### CAVE RUN LAKE 8,270 acres (Bath, Menifee, Rowan and Morgan counties)

Baring spring high water conditions, good growth of aquatic weed beds should occur; historically, Zilpo flats are covered with weed growth and fishing has been good in these areas for largemouth bass and muskellunge.

Bluegill	Good	Widespread within the lake, most 7 inches long or less, but it is not unusual to catch a few larger fish. Bluegill can be found in the back of coves or other quickly warming shallow water areas in the spring and on department placed or natural brush piles later in the year.
Catfish	Good	Good numbers of channel catfish up to 24 inches are fairly common along with a few larger sizes. Many are found along shallow rocky shoreline areas in early May-June where they seek cover in rock piles and submerged logs to spawn. Later they tend to be found in deeper water. Although not as plentiful, some trophy sized flathead catfish are found in the lake.

Location/Species	Forecast	Comments
CAVE RUN LAKE (co	ntinued)	
Crappie	Excellent	In recent years anglers have been experiencing outstanding crappie fishing. High numbers of fish in the 2 to 3 year old range in angler creels from 2015 will translate to another good year in 2016. Starting in May, fish around department placed and natural brush piles. If there are weeds present, summer fishing will be tougher; fish are still there just not as congregated around brush piles.
Largemouth bass	Good	Fish are heavy and healthy for their size with good numbers of larger fish (over the 16-inch upper slot limit) for a low nutrient reservoir. Try fishing the rocky banks warmed by the morning sun in April and May. Summer success is dependent on weed cover; fish the edges and pockets of the weed beds. If spring flooding limits weed growth, be prepared to search for summertime fish. <b>Note: 13-16 inch protective slot limit.</b>
Muskellunge	Excellent	Excellent angling continues with this fishery. Trophy fish can be found primarily in the spawning month of April in the backs of the large coves or in the fall months of September and October in association with gentle sloping gravely banks or near channel drop offs. <b>Note: 36 inch minimum size, one fish creel limit.</b>
Smallmouth bass	Fair	While still uncommon, smallmouth have been picking up in recent years. They are found primarily off rocky points or shoreline in the lower lake. <b>Note: 18-inch minimum size limit.</b>
Spotted bass	Good	Numerous within the middle and lower main lake shoreline; become less common within the upper reaches of the lake. Most less than 13 inches long.
White bass	Poor	This population is continuing to rebound from the 2013 die off. High numbers of age-0 to age-1 fish were observed in electrofishing studies in 2015, these fish will push through to larger sizes in 2016. Watch for fish in the jumps in the main lake area during the summer. Note: no more than 5 white bass in the daily creel limit may be greater than 15-inches long.
CEDAR CREEK LAKI	E 784 acres	(Lincoln County)
Bluegill	Good	(Lincoln County)  Numerous, although most are less than 7 inches. Bank anglers can choose from several designated fishing access areas for family fishing.
Channel catfish	Fair	Fair numbers of 12-18 inch fish with a few over 20 inches. Stocked every other year.
Crappie	Good	Stable crappie population provides good numbers and a fair number of fish over 10 inches. Try jigs tipped with minnows around submerged brush, particularly in late fall.
Largemouth bass	Excellent	Wide range of sizes available for anglers, with a high density of 15-19-inch fish. Although the numbers of fish >20 inches seems to have plateaued in recent years, Cedar Creek Lake remains a great place to catch a large bass. <b>Note: a trophy regulation of a 20-inch minimum size limit, 1 fish creel limit, is in effect on this lake.</b>
Rainbow trout	Fair	Stocked in late fall and early spring to provide cool weather fishery.
Redear sunfish	Good	Less numerous than bluegill but good numbers of 6-8 inch fish with a few over 10 inches available. Try mealworms around vegetation for best fishing.

Location/Species	Forecast	Comments
<b>CLEAR CREEK LAKE</b>	40 acres	(Bath County)
Bluegill	Good	Good numbers of $6-7$ inch bluegill are present with occasional 8 inch fish caught. Fish shallow shoreline areas in the spring especially behind standing timber.
Largemouth bass	Fair	Fair population with a few fish over 20 inches in the samples. Anglers should focus around the dam and the standing timber at the back of the lake.
Redear sunfish	Good	Good population of $7-8$ inch fish. Fish shallow shoreline areas in the spring especially behind standing timber.
CORINTH LAKE 96	acres (Gra	nt County)
Bluegill	Good	Good numbers of 6-8 inch fish; >8 inch fish present. Numbers of quality-size fish continue to increase from shad eradication.
Channel catfish	Good	Catchable-size channel catfish stocked in 2015; fish over 15 inches common.
Crappie	Fair	Low numbers; most fish between 8 to 9 inches.
Largemouth bass	Fair/Good	Good numbers of bass at and above size limit (12-inch size limit); excellent potential for bass between 15-20 inches.
Redear sunfish	Good	Increasing numbers of fish greater than 7 inches; numbers of quality-size fish continue to increase from shad eradication.
Bluegill	Good	Bluegill are fairly abundant and some are excellent size (>8 inches). Try crickets in coves and cuts for good early summer action or spinner rigs with half nightcrawlers or
Channel catfish	Excellent	Excellent population of 2-5 pound channels with some larger fish in the mix. Nightcrawlers still fished or slowly trolled on main lake flats and cuts during late summer or fall should pick up some quality fish.
Crappie	Excellent	Modest density, but very good size range. Large crappie (12-14 inches) fairly abundant compared to historical averages. The flooded trees should benefit the crappie population for the foreseeable future. Try jigs or minnows in headwaters of the lake (near mouth of Laurel and Rockcastle Rivers) in early spring for some excellent fishing. Fall fishing should also be good in major tributaries.
Largemouth bass	Good	Good density of fish up to 18 inches; some larger fish are available. Creek arms in upper lake are slightly better for largemouth than lower lake arms.
Smallmouth bass	Good	Good numbers of 14-18 inch fish present with moderate density of 18-inch plus fish. Try jigs or float-n-fly techniques for good winter fishing.
Spotted bass	Good	Good density of 12-15 inch fish; some larger ones are available. Found throughout lake, although more common in lower half.
Striped bass	Good	The striped bass population continues to improve and anglers can expect to catch slightly larger fish in 2016. Multiple year classes are contributing to the fishery and increasing numbers of fish are > 30 inches this year. The 22-inch size limit remains in effect, and anglers should have no trouble catching their limit.

Location/Species	Forecast	Comments
CUMBERLAND LAKE	E (continued)	
Walleye	Good	Walleye numbers should continue to improve in 2016. The population still lacks large fish but anglers should find adequate numbers of 2-4 pounders. Best fishing in spring is at night on surface baits. Night crawler rigs and deep diving crank baits trolled 20-35 feet deep along main lake shoreline work well during summer and fall.
White bass	Fair	Population level remains low but size is good. Upper half of lake is best for numbers. For best success, try headwater areas in early spring.
CUMBERLAND LAKE	E TAILWATER	(Russell, Cumberland, Clinton, and Monroe counties)
The new lower sec	tion of Hatchery	Creek will open in Spring 2016. Check the KDFWR website (fw.ky.gov) for updates on the ion will be catch-and-release only.
Brook trout	Fair	Annual stockings continue in the tailwater. Most fish range from 10-14 inches, but some larger fish are present. The state record remains vulnerable in 2016 as brooks grow into the larger size ranges. Brook trout are aggressive feeders that bite readily on a variety of baits.
Brown trout	Good	Anglers can expect good numbers of browns up to 16 inches, and fair numbers of large fish. The abundance of keeper fish (>20 inches) remains low but should improve in the coming years.
Rainbow trout	Good	Rainbow trout abundance is still lower than peak years, but adequate numbers of stocked fish should be available to anglers. Anglers should find pretty good numbers of fish in the slot (15-20 inch), but numbers of fish over 20 inches is still down.
Sauger	Fair	Population tends to fluctuate yearly. Try near dam in late winter and early spring with small jigs and grubs. From mid-summer to fall, fish below Burkesville trolling shad colored crankbaits or nightcrawler rigs.
Striped bass	Fair/Good	Fairly low density population but good size (20 lbs +). The tailwater is the best bet for a trophy striper in the state. Fishing best from Burkesville area downstream during summer months using live bait or trolling shad colored crankbaits or swim baits. Fish can be found throughout the tailwater during other seasons.
Walleye	Fair	Moderate numbers but good size (2-5 lb). Fish dam area in late February – mid March. From mid-summer to fall, fish below Burkesville trolling shad colored crankbaits or nightcrawler rigs.
White bass	Good	Best in lower sections near creek mouths during late winter – early spring.
DALE HOLLOW LAK		s in Kentucky (Clinton and Cumberland counties)
Largemouth bass	Good	Good numbers of larger (16-19 inch) fish.
Smallmouth bass	Good	Consistent fishery continues to produce impressive numbers of fish over 3 lbs. Standard fare of jigs and soft plastics work well as do float-n-fly techniques in cold water months.
Spotted bass	Good	Good density of quality-sized fish (12-15 inches).

Location/Species	Forecast	Comments
DEWEY LAKE 1,10	00 acres (Flo	yd County)
		sent at Dewey Lake. Boaters are asked to take measures to prevent the spread of these cies by properly inspecting and disinfecting boats, trailers and equipment.
Blue catfish	Good	Jugging produced good numbers of fish during summer 2015. Expect same pattern in 2016. Larger fish are approximately 36 inches.
Bluegill	Fair	Stable population with good growth. Large fish are around 7 inches long.
Channel catfish	Excellent	Good numbers found in the lower lake during winter months. The upper lake is best in the summer months. Lots of good bank fishing access.
Crappie	Good	Black and white crappie are present with larger white crappie to 16 inches and black crappie to 11 inches. Popular with bank anglers.
Flathead catfish	Good	Good size distribution; a few large fish from 20-40 pounds caught each year. Popular with ticklers/noodlers in June; try fishing shallow in September and October.
Largemouth bass	Good	Improving numbers of fish 15-20 inches. March is good time to go after largest fish in shallow murky water with artificial plastics. When water is clear fish group close to submerged brush and trees.
Muskellenge	N/A	Stocked for first time in October 2014. Fall 2015 seeing some 24-25 inch fish. Annual stockings will continue when excess hatchery fish are available. <b>Note: March 1, 2016 new minimum size regulation of 36 inches.</b>
Redear sunfish	Fair	Fish are present through 12 inches. Some areas such as the rip-rap rock face of dam hold good fish. Increasing population of zebra mussels will improve this fishery. Try fishing bait near the bottom around weedy areas or brush. If catching bluegill, then go deeper or further off shoreline to get into redear sunfish.
White bass	Fair	Numbers remain stable. John's Creek will congregate fish in March-April for spawning (look for Redbud trees in bloom). Later in September look for good numbers of fish in shallow water areas in head of coves off main lake after shad.
DIX RIVER - below He	errington Lake (	Garrard and Mercer counties)
Brown Trout	Fair/Good	Low numbers of holdover fish are expected due to flooding rains in spring of 2015. One stocking (usually June) per year.
Rainbow Trout	Fair/Good	Low numbers of holdover fish are expected due to poor water conditions in fall of 2015. Stocked on a monthly basis, 9 out of 12 months. Use of artificial lures or flies only from Herrington Lake Dam downstream for 2 miles.
FI KHORN CREEK -	main stem and l	North Fork (Franklin and Scott counties)
Catfish	Good	Large channel catfish possible in bigger pools; good numbers of "keeper" channel catfish, especially in the lower section of the main stem. Flathead catfish are present.
Largemouth bass	Good	Quality fishing found in slow moving pools in the North Fork. <b>Note: 12-16 inch slot limit applies on main stem.</b>
Rock Bass	Good	Many fish between 7-9 inches in main stem and in the lower North Fork section.
Smallmouth bass	Excellent	Very good numbers of smallmouth bass, especially on the main stem within the slot. Also good numbers of fish greater than 16 inches. <b>Note: fish from 12-16 inches (slot limit) are protected in the main stem Elkhorn Creek.</b>

Location/Species	Forecast	Comments
ELMER DAVIS LAKE	149 acres (	Owen County)
Bluegill	Fair	Good numbers of fish between 6-8 inches; many found around beaver lodges and boat docks. Bluegill population is being negatively affected by presence of gizzard shad in the lake.
Channel catfish	Good	Catchable-size fish stocked last year; fish over 15 inches possible.
Crappie	Fair/Good	Most around 9-11 inches.
Largemouth bass	Good/Excellent	Plenty of bass under 12-15 inch protected slot limit with good numbers in the slot. Good numbers of bass greater than 15 inches. Weed beds are back. Trophy-size bass are possible.
Redear sunfish	Good	Good numbers of fish between 6-8 inches near weed beds or on spawning beds in upper parts of embayments. "Shellcrackers" greater than 10 inches are present. Population is being negatively affected by shad.
Warmouth	Fair	Nice-size fish present, usually around weed beds. Locally known as "redeye" or "rockbass".

#### FAGAN BRANCH LAKE 140 acres (Marion County)

Summertime (mid-June through September) temperature and oxygen levels for trout in the 20-30 foot range. This zone decreases as the summer progresses. Lake is extremely clear year-round.

Bluegill	Fair	Fair numbers of 7-inch plus fish available.
Channel catfish	Good	Good numbers of 11-14 inch fish. Stocked every other year.
Crappie	Fair	Best fishing early and late and at night under floating lights using minnows. Fair numbers of fish, but not many larger fish (10-inch plus).
Largemouth bass	Fair	Stable population but slower growing. Fair numbers of 12-inch plus fish available.
Rainbow trout	Fair/Good	Best fishing will be at night under floating lights from May–October due to clear water. Fish will be in 20-30 foot ranges during the summer. Fish available throughout the lake during the winter. Stocked semi-annually (November and February) since 2010. 15-inch plus fish available.
Redear sunfish	Good/Excellent	Above average numbers of larger fish (8-inch plus) present. Fish edges of deep weed beds (15-20 feet) during summer.

#### FISHTRAP LAKE and TAILWATER 1,131 acres (Pike County)

Zebra mussels are present at Fishtrap Lake. Boaters are asked to take measures to prevent the spread of this damaging non-native invasive species by properly inspecting and disinfecting boats, trailers and equipment.

Blue catfish	Good	Fish were stocked again in 2015 and look for numbers to remain stable. Primarily feed on zebra mussels in spring and fall and on gizzard shad during summer and winter. Blues were first stocked in 2011 and larger fish are approximately 32-34 inches.
Bluegill	Excellent	Some large bluegill available up to 11-12 inches; May and June have been popular months for angling.
Channel catfish	Excellent	Good size distribution and numbers found throughout the lake and into Levisa Fork.  Provide good opportunity for tickling/noodling during nesting.

Location/Species	Forecast	Comments
FISHTRAP LAKE and	TAILWATER (c	ontinued)
Crappie	Excellent	Distribution and numbers are good through 13 inches with some larger fish available to 15-16 inches.
Flathead catfish	Excellent	Numerous large fish with good numbers found in the lake and into Levisa Fork. Numerous rocky ledges and banks around the lake provide excellent noodling opportunities when fish are spawning.
Hybrid striped bass	Excellent	Very good numbers of hybrids in the lower lake in late summer and at winter pool; fair spawning run in headwaters in April. Night fishing has become popular; look for fish in the jumps from May-July; good distribution of fish through 10 pounds.
Largemouth bass	Good	Fair numbers with excellent size distribution and good growth; lake is prone to rapid changes in water level affecting fish activity for periods. Ratio of keeper size bass is one of the best in eastern Kentucky.
Redear sunfish	Fair	This fishery was initiated with stockings from 2010-2013. Larger fish are approximately 10-12 inches.
Smallmouth bass	Good	Make up approximately 15% of black bass; good fishing primarily in the headwaters of the lake up into the Levisa Fork River and in the lower portion of the lake from Grapevine to dam. Large individuals up to 22 inches. Excellent stream fishing in the Levisa Fork upstream of the lake.

GRAYSON LAKE	1,512 acres	(Carter and Elliott counties)
Bluegill	Good	Numerous, up to 7 inches long. Widespread throughout the lake as well as other kinds of panfish. Try fishing 5-6 feet deep in May and June.
Catfish	Good	Widespread throughout the lake. Fair numbers of channel and flathead catfish up to 20 inches long, some trophy-size flathead catfish. Try fishing the middle and upper sections in early fall.
Crappie	Fair	Population appears to be on the upswing. Search out cover (there are several department-placed brush piles placed throughout the lake) or fish deep near channel edges around standing timber.
Hybrid striped bass	s Good	This fishery continues to improve beyond expectations. Several larger fish remain from previous stocking efforts and currently stocked fish are growing well. Fish are congregated in the Bruin area in the late fall.
Largemouth bass	Good	Fair numbers of fish over 15 inches, but most still less than the minimum size limit. Most prevalent of all black bass species in the lake. Best population exists in the Bruin areas. High water helps to congregate these fish within the flooded vegetation in the spring and the fall months and can produce feeding frenzies and high catch rates within the open water areas of Deer Creek and Bruin.  Note: 15 inch minimum size limit.
Spotted bass	Fair	Stable population. Most fish are found in lower 2/3 of lake.
Smallmouth bass	Poor	Rarely seen while sampling. The best bet is the lower portion of the lake, specifically around the Deer Creek area near the dam. <b>Note: 15 inch minimum size limit.</b>

Location/Species	Forecast	Comments
		own to mouth at Spottsville)
Catfish	Good	Good populations of channel, blue, and flathead catfish. Trophy-size blue and flathead catfish available. Good bets are tailwaters below locks and dams, in deeper holes, and around logs, brush, or rocks.
Crappie	Fair/Good	Fish tributary mouths, below locks and dams, and brush along bank in spring and fall with live minnows.
Striped and hybri striped bass	d Good	Stripers and hybrids congregate below dams late spring through early fall. Cast top water plugs or lead head jigs with plastic bodies into boils below dams. Shad imitation swim baits and bucktail jigs also produce well.
White bass	Good/Excellent	Fish mouth of tributary streams in spring and below dams mid-spring through fall. Top water baits or lead head jigs below dams and lead head jigs or live bait at mouth of tributary streams during spring. Fish bank access area and boat ramp at mouth of Pond River late March through April.
GREEN RIVER (	dam at Rochester un	o to Green River Lake dam)
Muskellunge	Good	Fishing best in spring at feeder creek mouths and around woody debris or other current breaks. Fishing below instream dams also good bet. Stocked annually throughout river above Rochester.
Rock bass	Excellent	Good numbers of 6-8 inch fish present from dam to Munfordville.
Smallmouth bass	Good	Good numbers overall with larger fish (15-inch plus) available. Fishing best from Green River Lake dam to Mammoth Cave.
Walleye/sauger	Fair/Good	Fishing best below Green River Lake dam area and instream dams from early winter through spring. Walleye most common of the two species.
GREEN RIVER LAK	<b>(E</b> 8,210 acres	(Taylor and Adair counties)
Summertime (mid	d-June through mid-0	October) oxygen levels are too low to support fish below 18-20 feet.
Bluegill	Fair/Good	Large population with occasional 7-8 inch fish. Fish stump beds and back ends of coves during spring and deep-water brush, standing timber and structure during summer for larger fish.
Channel catfish	Good/Excellent	All sizes available. Hook and line fishing best at night or when water levels are on the rise and muddy. Jug fishing good mid-summer to fall.
Largemouth bass	Good/Excellent	Good numbers of quality-sized fish (15-inch plus) and better than average numbers of 18-inch plus fish.
Muskellunge	Good	Legal-sized fish (36-inch) and trophy fish (40-inch plus) numbers fair. Fall, late winter and spring excellent time to catch trophy fish. Population maintained by annual stocking.
Smallmouth bass	s Fair	Low density overall; however, can expect a slight upturn in numbers from good spawn of 2010. Expect better than average numbers of 15-inch plus fish. Fish in mid-lower sections of Robinson Creek and Green River arms for best numbers and chance for 20-inch plus fish.
Spotted bass	Good	Better than average numbers of 12-inch plus fish. 15-inch plus spots available.

Location/Species	Forecast	Comments
<b>GREEN RIVER LAKE</b>	(continued)	
Walleye	Fair	Overall numbers are fair as are numbers of larger fish (20-inch plus). Fish dam area and headwater areas of Robinson Creek and Green River arms in late February through mid to late March. Target standing timber coves, main lake stump beds, humps, and channel drop offs during summer and fall. Population maintained by annual stocking.
White crappie	Good/Excellent	Good numbers of harvestable-size fish (9-inch plus). Expect to weed through lots of sub-legal fish to find keeper fish. Keep an eye out for the occasional black crappie.
GREENBO LAKE 1	81 acres (Gree	enup County - Greenbo State Park)
Bluegill	Fair	Increases in the growth of shoreline and aquatic vegetation will continue to help this population. Fish deep water structure in the spring and beaver lodges in the summer.  Note: bluegill and other sunfish - 15 fish daily creel limit.
Channel catfish	Good	Stocked every other year; many 2-3 pound fish observed during fall sampling. <b>Note: limb lines (set lines), jug lines, and trot lines are illegal to use at this lake.</b>
Largemouth bass	Good	Good numbers of fish over 15 inches long and chances for trophy sized fish over 20 inches. Due to water clarity, anglers may have to fish deeper or at night. Fish around downed trees or brush piles.
Rainbow trout	Excellent	Annually stocked population with some carry-over; best fishing year-round is in the dam area; during the spring and summer try fishing at night under lights. Winter fishing can also be good.
Redear sunfish	Fair	Numbers low; however a few over 8 inches can be caught. After the spawn try fishing in deeper water. <b>Note: see bluegill regulation.</b>
GUIST CREEK LAKE	317 acres (	Shelby County)
Bluegill	Fair	Abundant; most less than 6 inches. Easy to catch; good place to take a child fishing with live bait (redworms or crickets).
Catfish	Excellent	Catchable-size channel catfish stocked last year; very good numbers of catfish with potential for trophies. Flatheads present in lake. <b>Note: this lake is home to current white catfish state record.</b>
Common carp	Good	Fish in coves or headwaters of the lake; fish with grain-based bait; 3 to 4 lb. fish common.
Crappie	Fair/Good	Most in the 7-10 inch range; larger fish are present.
Hybrid striped bass	Fair	Small population; good numbers of fish in the 16 to 22 inch range; potential for trophysized fish.
Largemouth bass	Good/Excellent	Very good numbers of fish over the 12-inch size limit. Excellent potential for quality-size fish and bass over 20 inches. Excellent potential for a trophy fish.
Saugeye	Fair	Third year of stocking was completed in 2015; saugeye up to the 20 inches are present. <b>Note: 15-inch size limit</b>

Location/Species	Forecast	Comments
HERRINGTON LAKE	2,500 acres	(Mercer, Boyle and Garrard counties)
Bluegill	Good	Stable number of fish up to 8 inches can be found around cover in embayments and inlets of rock walls. Best April-June.
Common carp	Fair	Fish in coves or headwaters with grain-based baits on bottom.
Crappie	Good	Difficult to locate but many quality-size fish (9 inches or larger) available; best around brush or fallen trees in upper half of lake. Potential for large black crappie around debris in inlets in main lake.
Flathead catfish	Good	All sizes present; usually feed at night on sunfish and shad.
Hybrid striped bass	Fair	Overall numbers are down, but 2 to 5-lb fish are present; possible trophy-sized fish.
Largemouth bass	Good/Excellent	Good numbers of fish over 12 inches, many over 15 inches; largest fish around shoreline cover in March and April; potential for trophy-sized fish. Best success in the upper half of lake. Good at night during the summer.
Longnose gar	Good	Large fish common; look for them around schools of shad. Schooling common in fall; potential trophy size.
White bass	Fair/Good	Numbers are down due to recent fish kill caused by disease, but fair numbers of fish in the 12-14 inch range; Trophy-size present.

#### KENTUCKY LAKE and TAILWATER 55,025 acres (Marshall, Calloway, Lyon, Livingston and Trigg counties)

night crawlers.

Asian carp (silver, bighead, and grass carp) are present in the lake and tailwaters. Although unwanted, this has created an excellent bowfishing opportunity and a way that anglers can help remove these unwanted fish from our waters. Bighead carp are among the largest fish you will find in north America. The best fishing will be in the tailwaters where Asian carp and other rough species often congregate in the spring and summer. Asian carp make excellent table fare despite their reputation as a "trash fish" and preparation instructions can be found on the KDFWR webpage (fw.ky.gov). Remember to be courteous to other anglers and dispose of fish carcasses properly. Harvest of Asian carp is encouraged!!

other anglers and di	spose of fish card	casses property. Harvest of Asian carp is encouraged:
Blue catfish	Excellent	In the lake, best fishing is along the main river channel. Fish congregate along the ledges of the channel when there is current, and scatter along the adjacent flats during periods of low flow. Drift using live shiners or cut shad/skipjack in areas where creek channels meet the main river channel and along the main river channel near bends. Use a depth finder to determine if the fish are near the bottom or suspended, and then fish appropriately. Other baits to try include: chicken hearts, night crawlers, grass hoppers or catalpa worms.
		In the tailwaters there are year-round fishing opportunities. The ideal times are in April and May, and October and November. The best bait option is using fresh bait (shad and skipjack herring) caught in the tailwaters. Though most anglers fish from a boat, two fishing piers offer great bank fishing opportunities.
Bluegill	Excellent	There are high numbers of 6-8 inch fish. The best fishing is in May, but bluegill can be caught year-round. Crickets and worms fished under a cork during the spawn are great baits. Later in summer try the stake beds or deep water (6-10 foot) brush piles. Catch

bluegill in a feeding frenzy during the summer months where mayflies are hatching along the shoreline. Also try deeper rocky points during the summer with pieces of

Location/Species	Forecast	Comments
KENTUCKY LAKE ar	nd TAILWATER	(continued)
Channel catfish	Excellent	In the lake, the best fishing occurs during the late spring along main lake chunky rock shoreline and rip-rap shoreline around the dam and marinas. Good fishing is available along the river channel during summer and on flats along the secondary channels. Jug fishing in the embayments is often a good way to catch a mess of channel catfish.
		In the tailwaters there are year-round fishing opportunities. However, blue catfish outnumber the channel catfish in the tailwater area. Two piers offer good bank fishing access points, while fishing from a boat might be best downstream around woody debris, stumps and creek mouths.
Crappie	Good	Anglers fishing deep are catching fair numbers of keeper-size white crappie. Anglers fishing shallow along rocky shoreline are catching fair numbers of black crappie. Try casting toward the shoreline with a curly tail jig. Crappie are vulnerable during late winter or early spring as they move toward shallow water areas for spawning. Fish are shallow in early to mid April around buttonball bushes, brush piles and stake beds. Crappie move to secondary channels, drops and flats near creek channels during the fall and winter, and are most frequently found near structure. A lot of short crappie being caught; just means better fishing to come.
Largemouth bass	Excellent	Several good year classes have resulted in quality fishing. During late winter and early spring, suspended fish can often be caught with jerkbaits and other shad imitations such as the Alabama rig. As spring progresses, fish move to wood structure along the shoreline. During the summer big fish are found along the main lake drops and secondary channels. Fishing the weed beds is good with some top water baits on cloudy, rainy days during the summer. Smaller fish can be found in the backs of embayments around brush, and chasing shad during late summer. During the fall, fishing around boat docks and rocky shorelines is good.
		There is some fine bass fishing in the tailwaters below KY Dam. Try fishing white spinnerbaits along the rip rap shoreline.
Redear sunfish	Excellent	Best fishing is in late April and early May. Redear like areas with gravel substrate. If there are yellow mustard flowers inundated by water in the backs of embayments during this time, this is a good place to find redear. Submerged pondweed clumps also hold redear sunfish. These fish primarily search the bottom for bugs, snails and small mussels; hence you need to fish near or on the bottom. Redear generally spawn a little deeper than bluegill.
Sauger	Fair	In the lake, there is a moderate population of harvestable-size (≥14 inch) fish. Best fishing is in fall and winter along secondary channels. Try trolling crankbaits.
		In the tailwaters there is a higher abundance of harvestable-size sauger. This is a good winter fishery. Boat fishing is ideal, though two new fishing piers offer good bank fishing opportunities.
Smallmouth bass	Good	Best fishing along rocky shorelines and bluffs, creek channels at mouths of embayments, the canal main lake points, and main lake stumps. Night fishing during the summer months along gravel bars at the mouths of LBL embayments and rocky shoreline is good with dark colored spinnerbaits with Colorado style blades or crayfish color creature baits and jigs.
Spotted bass	Poor	Spotted bass are not normally targeted, but are caught while fishing for largemouth or smallmouth bass. Spotted bass migrate to the streams at the backs of embayments during the early spring. Gravel bars and points are prime areas to fish during the summer. Secondary points are good during the fall.

Location/Species	Forecast	Comments
KENTUCKY LAKE a	nd TAILWATER	(continued)
Striped bass	Good	Striped bass and its hybrid are being caught from the lake while fishing for white bass. Trolling with deep diver baits which imitate minnows is effective.
		Better fishing is available below Kentucky Dam in the tailwater. Fish congregate below the dam during periods of discharge through the generator section of dam. Live herring are the preferred bait. Fishing from a boat is best, but the best shoreline fishing would be off the two fishing piers near the dam. From the shoreline fish with 4-inch sassy shad type baits.
White bass	Fair	Fishing is good in the main lake during summer months when white bass are found schooling along secondary creek channels near the main river channel. Trolling can be very effective for locating these fish. Also look for jumps; when groups of white bass chase shad near the water's surface. In-line spinners and silver spoons work great for catching white bass. Look for main channel points and bends; also near the canal.
		Good numbers are also found in the Kentucky Dam tailwater area along with its cousin the yellow bass. When fishing from the bank or boat, try using small curly tail jigs or inline spinner type baits.
Yellow bass	Excellent	Lots of fish but they don't get very big. Their taste is much better than most anglers perceive. Casting the rocky shoreline with a curly tail jig or rooster tail works great. Large numbers are caught while casting the rocky shoreline in the spring for crappie.

KENTUCKY RIVER	255 miles (main	n stem)
Bluegill*	Fair	Shoreline areas around woody debris out of the current hold some quality fish but in low numbers.
Channel catfish	Fair	Look for silt-free rocky habitat or below dams or palisades.
Common carp	Fair	Bottom fish with grain-based baits below dams.
Crappie*	Fair	Currently good number of crappie in the Kentucky River; locate in backwater/creek-mouth areas with low current. Fish up to 12 inches or larger possible.
Flathead catfish	Fair	Large fish present around large boulders and log jams.
Hybrid striped bass	Fair	Sporadic catches of 15 to 20-inch fish below dams. Stocked annually.
Largemouth bass*	Fair	Locate in cover just below dams, in creek mouths, bank slides, or around timber. Numbers have been reduced due to recent flood events over the past year.
Muskellunge	Good	Best in spring around creek mouths, and in summer around timber and creek mouths. New study looking at lower-river stockings. Stocked annually in upper river. More fish in upper half of river (pools 6-14).
Sauger	Good	Found around sand bars, creek mouths, below dams around rock and sand bars.
Smallmouth bass	Fair	Nice fish possible in mouths of quality tributaries. Also found around rock piles and walls.
Spotted bass	Fair	Sporadic catches of small to medium-size fish; usually below dams but also around palisade rock wall areas.
Walleye	Fair	Present in low numbers due to past stockings and escapement from reservoirs in the headwaters; good-size fish possible below dams.

Forecast Comments	Comments	Forecast	Location/Species
ontinued)		(continued)	KENTUCKY RIVER (
ontinuou)		(oontinaoa)	REITIOOR RIVERY

White bass Good Many 10-12 inch fish below dams.

<sup>\*</sup>These are considered "standing water species" and do best in the river after prolonged periods of low-flow conditions. High water or flooding conditions for extended periods will reduce their numbers in the Kentucky River.

KINCAID LAKE	183 acres (F	Pendleton County)
Bluegill	Poor	Abundant, most fish are less than 6 inches.
Channel catfish	Good	Catchable-size stocked last year; good potential for fish over 15 inches.
Crappie	Fair	Abundant; most between 6-9 inches.
Largemouth bas	s Good/Exce	ellent Very good numbers of fish over the 12-inch size limit; good numbers of fish over 15 inches; excellent potential for a trophy fish.

#### KINGFISHER LAKES 13 acres and 21 acres (Daviess County)

Renovation work complete. The lakes are re-filling and were stocked with channel catfish, remedial bluegill, and remedial bass in the fall of 2015. Sampling to monitor the population will begin in spring 2017.

LAKE CARNICO	114 acres	(Nicholas County)
Bluegill	Poor	Numbers are low. Concentrate efforts around the shallow spawning beds in the backs of the coves in spring and around structure or weed beds in the summer.
Catfish	Fair	Fair numbers of flathead and channel catfish can be found. Lake is stocked with channel catfish.
Largemouth bas	ss Fair	Fair numbers of bass present, a few over 15 and 20 inches. Anglers should concentrate summer-time efforts around aquatic weed beds.
Redear sunfish	Poor	Numbers are low. See bluegill.

LAKE MALONE	767 acres (Muh	lenburg, Todd, and Logan counties)
Bluegill	Fair	Population mostly comprised of 4-6 inch fish.
Channel catfish	Good	Stocked annually, good number of 14-16 inch fish with a few 20-inch plus available.
Crappie	Fair/Good	Moderate number of 9-10 inch fish. Fish brush piles and fallen trees along some of the steeper banks.
Largemouth bass	s Excellent	Stable population with large number of 15-20 inch fish. Many trophy fish available. Fish the shoreline waterwillow during spring and fall and at night during summer with plastic baits. Also try submerged trees and brush along deeper banks in spring and summer with spinner baits and jigs.
Redear sunfish	Fair/Good	Low number of redear, but good size (9-11 inch) available. Fish water willow edge along the shoreline in spring-early summer. Move to deeper submerged structure as water warms during summer.

Channel catifish Good Stocked every other year. Sampling shows high numbers of these stocked-size fish and several over this size. Note: 12-inch minimum size limit; limb lines (set lines jug lines, and trot lines are illegal to use at this lake.  Largemouth bass Good Good numbers of bass over 15 and 20 inches present. Gibson Bay, upper end of th lake and away from the bank anglers are good areas during the summer.  Note: 15-inch minimum size limit.  Redear sunfish Fair Good numbers of fish in the 8 - 10 inch range; see bluegill.  AKE WILGREEN 169 acres (Madison County) Blue catfish Good Stocked every third year. Good catches regularly reported in the summer.  Bluegill Good Similar number as previous years and most fish are less than 6 inches. The populatis healthy up to that point.  Largemouth bass Excellent High numbers of fish over 15 and 20 inches. These sizes of fish are fat and healthy: their size. All indicators are continuing to show an increase in this population. Shou be another great year.  Redear sunfish Good Similar numbers to previous years and most fish are less than 7 inches; see bluegil  AUREL RIVER LAKE 5,830 acres (Whitley and Laurel counties) Bluegill Good Good summer fishery with fairly abundant 6-7 inch fish. Try crickets along rock walls upper Laurel River Arm during summer.  Crappie Fair Not a numbers game, but some larger (9-inch plus) crappie are caught each year. Upper Laurel River arm near city dam is best area.  Largemouth bass Good The largemouth population continues to improve. Currently there are good numbers 14-18-inch fish with some over 20 inches. Upper Laurel River arm is best area.  Rainbow trout Fair Stocked annually in January. Should be adequate numbers of 10-14 inch trout and few larger fish. Try trolling small cranks or spoons during winter and early spring for good catches of stocked fish and the occasional larger fish.		Forecast	Comments
Channel catfish Good Stocked every other year. Sampling shows high numbers of these stocked-size fish and several over this size. Note: 12-inch minimum size limit; limb lines (set line: jug lines, and trot lines are illegal to use at this lake.  Largemouth bass Good Good numbers of bass over 15 and 20 inches present. Gibson Bay, upper end of the lake and away from the bank anglers are good areas during the summer.  Note: 15-inch minimum size limit.  Redear sunfish Fair Good numbers of fish in the 8 - 10 inch range; see bluegill.  AKE WILGREEN 169 acres (Madison County) Blue catfish Good Stocked every third year. Good catches regularly reported in the summer.  Bluegill Good Similar number as previous years and most fish are less than 6 inches. The populat is healthy up to that point.  Largemouth bass Excellent High numbers of fish over 15 and 20 inches. These sizes of fish are fat and healthy their size. All indicators are continuing to show an increase in this population. Shou be another great year.  Redear sunfish Good Similar numbers to previous years and most fish are less than 7 inches; see bluegil  AUREL RIVER LAKE 5.830 acres (Whittley and Laurel counties) Bluegill Good Good summer fishery with fatrly abundant 6-7 inch fish. Try crickets along rock wall upper Laurel River Arm during summer.  Crappie Fair Not a numbers game, but some larger (9-inch plus) crappie are caught each year. Upper Laurel River Arm near city dam is best area.  Rainbow trout Fair Stocked annually in January. Should be adequate numbers of 10-14 inch trout and few larger fish. Try trolling small cranks or spoons during winter and early spring for good catches of stocked fish and the occasional larger fish.  Smallmouth bass Good Moderate density of 12-14 inch fish with a few exceeding 15 inches. Best in the mid and lower sections of the lake. Look for schooling fish chasing shad on the surface early fall.  Walleye Good Moderate density of 12-14 inch fish with a few exceeding 15 inches. Best in the mid and lower sections of the lake. Lo	AKE REBA 76 ac	res (Madisor	n County)
and several over this size. Note: 12-inch minimum size limit; limb lines (set line: jug lines, and trot lines are illegal to use at this lake.  Largemouth bass Good Good numbers of bass over 15 and 20 inches present. Gibson Bay, upper end of the lake and away from the bank anglers are good areas during the summer.  Note: 15-inch minimum size limit.  Redear sunfish Fair Good numbers of fish in the 8 - 10 inch range; see bluegill.  AKE WILGREEN 169 acres (Madison County)  Blue catfish Good Stocked every third year. Good catches regularly reported in the summer.  Bluegill Good Similar number as previous years and most fish are less than 6 inches. The populatis healthy up to that point.  Largemouth bass Excellent High numbers of fish over 15 and 20 inches. These sizes of fish are fat and healthy their size. All indicators are continuing to show an increase in this population. Shoube another great year.  Redear sunfish Good Similar numbers to previous years and most fish are less than 7 inches; see bluegil Good summer fishery with fairly abundant 6-7 inch fish. Try crickets along rock wall upper Laurel River Arm during summer.  Crappie Fair Not a numbers game, but some larger (9-inch plus) crappie are caught each year. Upper Laurel River Arm during summer.  Largemouth bass Good The largemouth population continues to improve. Currently there are good numbers 14-18-inch fish with some over 20 inches. Upper Laurel River arm is best area.  Rainbow trout Fair Stocked annually in January. Should be adequate numbers of 10-14 inch trout and few larger fish. Try trolling small cranks or spoons during winter and early spring for good actaches of stocked fish and the occasional larger fish.  Smallmouth bass Good Moderate density of 12-14 inch fish with a few exceeding 15 inches. Best in the mid and lower sections of the lake. Look for schooling fish chasing shad on the surface early fall.  Walleye Good Multiple year classes contributing to the fishery with good density of 20-23 inch fish surface baits at right during spring, o	Bluegill	Good	collected in 2015; most catchable fish are over 6 - 8 inches. Great lake to introduce
lake and away from the bank anglers are good areas during the summer.  Note: 15-inch minimum size limit.  Redear sunfish Fair Good numbers of fish in the 8 - 10 inch range; see bluegill.  AKE WILGREEN 169 acres (Madison County)  Blue catfish Good Stocked every third year. Good catches regularly reported in the summer.  Bluegill Good Similar number as previous years and most fish are less than 6 inches. The populat is healthy up to that point.  Largemouth bass Excellent High numbers of fish over 15 and 20 inches. These sizes of fish are fat and healthy their size. All indicators are continuing to show an increase in this population. Shou be another great year.  Redear sunfish Good Similar numbers to previous years and most fish are less than 7 inches; see bluegil Good Good summer fishery with fairly abundant 6-7 inch fish. Try crickets along rock wall: upper Laurel River Arm during summer.  Crappie Fair Not a numbers game, but some larger (9-inch plus) crappie are caught each year. Upper Laurel River arm near city dam is best area.  Largemouth bass Good The largemouth population continues to improve. Currently there are good numbers 14-18-inch fish with some over 20 inches. Upper Laurel River arm is best area.  Rainbow trout Fair Stocked annually in January. Should be adequate numbers of 10-14 inch trout and few larger fish. Try trolling small cranks or spoons during winter and early spring for good catches of stocked fish and the occasional larger fish.  Smallmouth bass Good Moderate density of 12-14 inch fish with a few exceeding 15 inches. Best in the mid and lower sections of the lake. Look for schooling fish chasing shad on the surface early fall.  Walleye Good Multiple year classes contributing to the fishery with good density of 20-23 inch fish. surface baits at night during spring, or troll deep diving crank baits or spinner rigs across points and islands in summer and fall.	Channel catfish	Good	Stocked every other year. Sampling shows high numbers of these stocked-size fish and several over this size. <b>Note: 12-inch minimum size limit; limb lines (set lines jug lines, and trot lines are illegal to use at this lake.</b>
AKE WILGREEN 169 acres (Madison County)  Blue catfish Good Stocked every third year. Good catches regularly reported in the summer.  Bluegill Good Similar number as previous years and most fish are less than 6 inches. The popular is healthy up to that point.  Largemouth bass Excellent High numbers of fish over 15 and 20 inches. These sizes of fish are fat and healthy their size. All indicators are continuing to show an increase in this population. Show be another great year.  Redear sunfish Good Similar numbers to previous years and most fish are less than 7 inches; see bluegil  Good Good summer fishery with fairly abundant 6-7 inch fish. Try crickets along rock wall upper Laurel River Arm during summer.  Crappie Fair Not a numbers game, but some larger (9-inch plus) crappie are caught each year. Upper Laurel River arm near city dam is best area.  Largemouth bass Good The largemouth population continues to improve. Currently there are good numbers 14-18-inch fish with some over 20 inches. Upper Laurel River arm is best area.  Rainbow trout Fair Stocked annually in January. Should be adequate numbers of 10-14 inch trout and few larger fish. Try troiling small cranks or spoons during winter and early spring for good catches of stocked fish and the occasional larger fish.  Smallmouth bass Good Moderate density of 12-14 inch fish with a few exceeding 15 inches. Best in the mid and lower sections of the lake. Look for schooling fish chasing shad on the surface early fall.  Walleye Good Multiple year classes contributing to the fishery with good density of 20-23 inch fish. surface baits at night during spring, or troll deep diving crank baits or spinner rigs across points and islands in summer and fall.	Largemouth bass	Good	
Bluegill Good Stocked every third year. Good catches regularly reported in the summer.  Bluegill Good Similar number as previous years and most fish are less than 6 inches. The populat is healthy up to that point.  Largemouth bass Excellent High numbers of fish over 15 and 20 inches. These sizes of fish are fat and healthy their size. All indicators are continuing to show an increase in this population. Show be another great year.  Redear sunfish Good Similar numbers to previous years and most fish are less than 7 inches; see bluegil Good Good summer fishery with fairly abundant 6-7 inch fish. Try crickets along rock wall upper Laurel River Arm during summer.  Crappie Fair Not a numbers game, but some larger (9-inch plus) crappie are caught each year. Upper Laurel River arm near city dam is best area.  Largemouth bass Good The largemouth population continues to improve. Currently there are good numbers 14-18-inch fish with some over 20 inches. Upper Laurel River arm is best area.  Rainbow trout Fair Stocked annually in January. Should be adequate numbers of 10-14 inch trout and few larger fish. Try trolling small cranks or spoons during winter and early spring for good catches of stocked fish and the occasional larger fish.  Smallmouth bass Good Persistent anglers can catch some very nice smallmouth at Laurel, including fish over pounds. Try casting jigs or swim baits in winter and early spring for best chance for trophy fish.  Spotted bass Good Moderate density of 12-14 inch fish with a few exceeding 15 inches. Best in the mid and lower sections of the lake. Look for schooling fish chasing shad on the surface early fall.  Walleye Good Multiple year classes contributing to the fishery with good density of 20-23 inch fish. surface baits at night during spring, or troll deep diving crank baits or spinner rigs across points and islands in summer and fall.	Redear sunfish	Fair	Good numbers of fish in the 8 - 10 inch range; see bluegill.
Bluegill Good Stocked every third year. Good catches regularly reported in the summer.  Bluegill Good Similar number as previous years and most fish are less than 6 inches. The populat is healthy up to that point.  Largemouth bass Excellent High numbers of fish over 15 and 20 inches. These sizes of fish are fat and healthy their size. All indicators are continuing to show an increase in this population. Show be another great year.  Redear sunfish Good Similar numbers to previous years and most fish are less than 7 inches; see bluegil Good Sood summer fishery with fairly abundant 6-7 inch fish. Try crickets along rock wall upper Laurel River Arm during summer.  Crappie Fair Not a numbers game, but some larger (9-inch plus) crappie are caught each year. Upper Laurel River arm near city dam is best area.  Largemouth bass Good The largemouth population continues to improve. Currently there are good numbers 14-18-inch fish with some over 20 inches. Upper Laurel River arm is best area.  Rainbow trout Fair Stocked annually in January. Should be adequate numbers of 10-14 inch trout and few larger fish. Try trolling small cranks or spoons during winter and early spring for good catches of stocked fish and the occasional larger fish.  Smallmouth bass Good Persistent anglers can catch some very nice smallmouth at Laurel, including fish over pounds. Try casting jigs or swim baits in winter and early spring for best chance for trophy fish.  Spotted bass Good Moderate density of 12-14 inch fish with a few exceeding 15 inches. Best in the mid and lower sections of the lake. Look for schooling fish chasing shad on the surface early fall.  Walleye Good Multiple year classes contributing to the fishery with good density of 20-23 inch fish. surface baits at night during spring, or troll deep diving crank baits or spinner rigs across points and islands in summer and fall.	AKE WII GREEN	169 acres (N	Madison County)
Bluegill Good Similar number as previous years and most fish are less than 6 inches. The popular is healthy up to that point.  Largemouth bass Excellent High numbers of fish over 15 and 20 inches. These sizes of fish are fat and healthy their size. All indicators are continuing to show an increase in this population. Shou be another great year.  Redear sunfish Good Similar numbers to previous years and most fish are less than 7 inches; see bluegil Good Good summer fishery with fairly abundant 6-7 inch fish. Try crickets along rock walls upper Laurel River Arm during summer.  Crappie Fair Not a numbers game, but some larger (9-inch plus) crappie are caught each year. Upper Laurel River arm near city dam is best area.  Largemouth bass Good The largemouth population continues to improve. Currently there are good numbers 14-18-inch fish with some over 20 inches. Upper Laurel River arm is best area.  Rainbow trout Fair Stocked annually in January. Should be adequate numbers of 10-14 inch trout and few larger fish. Try trolling small cranks or spoons during winter and early spring for good catches of stocked fish and the occasional larger fish.  Smallmouth bass Good Persistent anglers can catch some very nice smallmouth at Laurel, including fish ow pounds. Try casting jigs or swim baits in winter and early spring for best chance for trophy fish.  Spotted bass Good Moderate density of 12-14 inch fish with a few exceeding 15 inches. Best in the mid and lower sections of the lake. Look for schooling fish chasing shad on the surface early fall.  Walleye Good Multiple year classes contributing to the fishery with good density of 20-23 inch fish. surface baits at night during spring, or troll deep diving crank baits or spinner rigs across points and islands in summer and fall.			
is healthy up to that point.  Largemouth bass Excellent High numbers of fish over 15 and 20 inches. These sizes of fish are fat and healthy their size. All indicators are continuing to show an increase in this population. Show be another great year.  Redear sunfish Good Similar numbers to previous years and most fish are less than 7 inches; see bluegil Good Good summer fishery with fairly abundant 6-7 inch fish. Try crickets along rock walls upper Laurel River Arm during summer.  Crappie Fair Not a numbers game, but some larger (9-inch plus) crappie are caught each year. Upper Laurel River arm near city dam is best area.  Largemouth bass Good The largemouth population continues to improve. Currently there are good numbers 14-18-inch fish with some over 20 inches. Upper Laurel River arm is best area.  Rainbow trout Fair Stocked annually in January. Should be adequate numbers of 10-14 inch trout and few larger fish. Try trolling small cranks or spoons during winter and early spring for good catches of stocked fish and the occasional larger fish.  Smallmouth bass Good Persistent anglers can catch some very nice smallmouth at Laurel, including fish over pounds. Try casting jigs or swim baits in winter and early spring for best chance for trophy fish.  Spotted bass Good Moderate density of 12-14 inch fish with a few exceeding 15 inches. Best in the mid and lower sections of the lake. Look for schooling fish chasing shad on the surface early fall.  Walleye Good Multiple year classes contributing to the fishery with good density of 20-23 inch fish. surface baits at night during spring, or troll deep diving crank baits or spinner rigs across points and islands in summer and fall.			, ,
their size. All indicators are continuing to show an increase in this population. Show be another great year.  Redear sunfish Good Similar numbers to previous years and most fish are less than 7 inches; see bluegil  AUREL RIVER LAKE 5,830 acres (Whitley and Laurel counties)  Bluegill Good Good summer fishery with fairly abundant 6-7 inch fish. Try crickets along rock walls upper Laurel River Arm during summer.  Crappie Fair Not a numbers game, but some larger (9-inch plus) crappie are caught each year. Upper Laurel River arm near city dam is best area.  Largemouth bass Good The largemouth population continues to improve. Currently there are good numbers 14-18-inch fish with some over 20 inches. Upper Laurel River arm is best area.  Rainbow trout Fair Stocked annually in January. Should be adequate numbers of 10-14 inch trout and few larger fish. Try trolling small cranks or spoons during winter and early spring for good catches of stocked fish and the occasional larger fish.  Smallmouth bass Good Persistent anglers can catch some very nice smallmouth at Laurel, including fish over pounds. Try casting jigs or swim baits in winter and early spring for best chance for trophy fish.  Spotted bass Good Moderate density of 12-14 inch fish with a few exceeding 15 inches. Best in the mid and lower sections of the lake. Look for schooling fish chasing shad on the surface early fall.  Walleye Good Multiple year classes contributing to the fishery with good density of 20-23 inch fish. surface baits at night during spring, or troll deep diving crank baits or spinner rigs across points and islands in summer and fall.	Bluegill	Good	Similar number as previous years and most fish are less than 6 inches. The popular is healthy up to that point.
AUREL RIVER LAKE 5,830 acres (Whitley and Laurel counties)  Bluegill Good Good summer fishery with fairly abundant 6-7 inch fish. Try crickets along rock walls upper Laurel River Arm during summer.  Crappie Fair Not a numbers game, but some larger (9-inch plus) crappie are caught each year. Upper Laurel River arm near city dam is best area.  Largemouth bass Good The largemouth population continues to improve. Currently there are good numbers 14-18-inch fish with some over 20 inches. Upper Laurel River arm is best area.  Rainbow trout Fair Stocked annually in January. Should be adequate numbers of 10-14 inch trout and few larger fish. Try trolling small cranks or spoons during winter and early spring for good catches of stocked fish and the occasional larger fish.  Smallmouth bass Good Persistent anglers can catch some very nice smallmouth at Laurel, including fish over pounds. Try casting jigs or swim baits in winter and early spring for best chance for trophy fish.  Spotted bass Good Moderate density of 12-14 inch fish with a few exceeding 15 inches. Best in the mid and lower sections of the lake. Look for schooling fish chasing shad on the surface early fall.  Walleye Good Multiple year classes contributing to the fishery with good density of 20-23 inch fish. surface baits at night during spring, or troll deep diving crank baits or spinner rigs across points and islands in summer and fall.	Largemouth bass	Excellent	High numbers of fish over 15 and 20 inches. These sizes of fish are fat and healthy their size. All indicators are continuing to show an increase in this population. Show be another great year.
AUREL RIVER LAKE 5,830 acres (Whitley and Laurel counties)  Bluegill Good Good summer fishery with fairly abundant 6-7 inch fish. Try crickets along rock walls upper Laurel River Arm during summer.  Crappie Fair Not a numbers game, but some larger (9-inch plus) crappie are caught each year. Upper Laurel River arm near city dam is best area.  Largemouth bass Good The largemouth population continues to improve. Currently there are good numbers 14-18-inch fish with some over 20 inches. Upper Laurel River arm is best area.  Rainbow trout Fair Stocked annually in January. Should be adequate numbers of 10-14 inch trout and few larger fish. Try trolling small cranks or spoons during winter and early spring for good catches of stocked fish and the occasional larger fish.  Smallmouth bass Good Persistent anglers can catch some very nice smallmouth at Laurel, including fish over pounds. Try casting jigs or swim baits in winter and early spring for best chance for trophy fish.  Spotted bass Good Moderate density of 12-14 inch fish with a few exceeding 15 inches. Best in the mid and lower sections of the lake. Look for schooling fish chasing shad on the surface early fall.  Walleye Good Multiple year classes contributing to the fishery with good density of 20-23 inch fish. surface baits at night during spring, or troll deep diving crank baits or spinner rigs across points and islands in summer and fall.	Redear sunfish	Good	Similar numbers to previous years and most fish are less than 7 inches: see bluegil
Crappie Fair Not a numbers game, but some larger (9-inch plus) crappie are caught each year. Upper Laurel River arm near city dam is best area.  Largemouth bass Good The largemouth population continues to improve. Currently there are good numbers 14-18-inch fish with some over 20 inches. Upper Laurel River arm is best area.  Rainbow trout Fair Stocked annually in January. Should be adequate numbers of 10-14 inch trout and few larger fish. Try trolling small cranks or spoons during winter and early spring for good catches of stocked fish and the occasional larger fish.  Smallmouth bass Good Persistent anglers can catch some very nice smallmouth at Laurel, including fish over pounds. Try casting jigs or swim baits in winter and early spring for best chance for trophy fish.  Spotted bass Good Moderate density of 12-14 inch fish with a few exceeding 15 inches. Best in the mid and lower sections of the lake. Look for schooling fish chasing shad on the surface early fall.  Walleye Good Multiple year classes contributing to the fishery with good density of 20-23 inch fish. surface baits at night during spring, or troll deep diving crank baits or spinner rigs across points and islands in summer and fall.			Good summer fishery with fairly abundant 6-7 inch fish. Try crickets along rock walls
Largemouth bass Good The largemouth population continues to improve. Currently there are good numbers 14-18-inch fish with some over 20 inches. Upper Laurel River arm is best area.  Rainbow trout Fair Stocked annually in January. Should be adequate numbers of 10-14 inch trout and few larger fish. Try trolling small cranks or spoons during winter and early spring for good catches of stocked fish and the occasional larger fish.  Smallmouth bass Good Persistent anglers can catch some very nice smallmouth at Laurel, including fish over pounds. Try casting jigs or swim baits in winter and early spring for best chance for trophy fish.  Spotted bass Good Moderate density of 12-14 inch fish with a few exceeding 15 inches. Best in the mid and lower sections of the lake. Look for schooling fish chasing shad on the surface early fall.  Walleye Good Multiple year classes contributing to the fishery with good density of 20-23 inch fish. surface baits at night during spring, or troll deep diving crank baits or spinner rigs across points and islands in summer and fall.	Crappie	Fair	
few larger fish. Try trolling small cranks or spoons during winter and early spring for good catches of stocked fish and the occasional larger fish.  Smallmouth bass Good Persistent anglers can catch some very nice smallmouth at Laurel, including fish over pounds. Try casting jigs or swim baits in winter and early spring for best chance for trophy fish.  Spotted bass Good Moderate density of 12-14 inch fish with a few exceeding 15 inches. Best in the mid and lower sections of the lake. Look for schooling fish chasing shad on the surface early fall.  Walleye Good Multiple year classes contributing to the fishery with good density of 20-23 inch fish. surface baits at night during spring, or troll deep diving crank baits or spinner rigs across points and islands in summer and fall.			
pounds. Try casting jigs or swim baits in winter and early spring for best chance for trophy fish.  Spotted bass Good Moderate density of 12-14 inch fish with a few exceeding 15 inches. Best in the mid and lower sections of the lake. Look for schooling fish chasing shad on the surface early fall.  Walleye Good Multiple year classes contributing to the fishery with good density of 20-23 inch fish. surface baits at night during spring, or troll deep diving crank baits or spinner rigs across points and islands in summer and fall.	Largemouth bass	Good	Upper Laurel River arm near city dam is best area.  The largemouth population continues to improve. Currently there are good numbers
and lower sections of the lake. Look for schooling fish chasing shad on the surface early fall.  Walleye  Good  Multiple year classes contributing to the fishery with good density of 20-23 inch fish. surface baits at night during spring, or troll deep diving crank baits or spinner rigs across points and islands in summer and fall.	-		Upper Laurel River arm near city dam is best area.  The largemouth population continues to improve. Currently there are good numbers 14-18-inch fish with some over 20 inches. Upper Laurel River arm is best area.  Stocked annually in January. Should be adequate numbers of 10-14 inch trout and few larger fish. Try trolling small cranks or spoons during winter and early spring for
surface baits at night during spring, or troll deep diving crank baits or spinner rigs across points and islands in summer and fall.	Rainbow trout	Fair	Upper Laurel River arm near city dam is best area.  The largemouth population continues to improve. Currently there are good numbers 14-18-inch fish with some over 20 inches. Upper Laurel River arm is best area.  Stocked annually in January. Should be adequate numbers of 10-14 inch trout and few larger fish. Try trolling small cranks or spoons during winter and early spring for good catches of stocked fish and the occasional larger fish.  Persistent anglers can catch some very nice smallmouth at Laurel, including fish over pounds. Try casting jigs or swim baits in winter and early spring for best chance for
White bass Poor Poor population overall but a few can be caught in the Laurel River Arm.	Rainbow trout  Smallmouth bass	Fair Good	Upper Laurel River arm near city dam is best area.  The largemouth population continues to improve. Currently there are good numbers 14-18-inch fish with some over 20 inches. Upper Laurel River arm is best area.  Stocked annually in January. Should be adequate numbers of 10-14 inch trout and few larger fish. Try trolling small cranks or spoons during winter and early spring for good catches of stocked fish and the occasional larger fish.  Persistent anglers can catch some very nice smallmouth at Laurel, including fish over pounds. Try casting jigs or swim baits in winter and early spring for best chance for trophy fish.  Moderate density of 12-14 inch fish with a few exceeding 15 inches. Best in the mid and lower sections of the lake. Look for schooling fish chasing shad on the surface
	Rainbow trout  Smallmouth bass  Spotted bass	Fair Good Good	Upper Laurel River arm near city dam is best area.  The largemouth population continues to improve. Currently there are good numbers 14-18-inch fish with some over 20 inches. Upper Laurel River arm is best area.  Stocked annually in January. Should be adequate numbers of 10-14 inch trout and few larger fish. Try trolling small cranks or spoons during winter and early spring for good catches of stocked fish and the occasional larger fish.  Persistent anglers can catch some very nice smallmouth at Laurel, including fish over pounds. Try casting jigs or swim baits in winter and early spring for best chance for trophy fish.  Moderate density of 12-14 inch fish with a few exceeding 15 inches. Best in the mid and lower sections of the lake. Look for schooling fish chasing shad on the surface early fall.  Multiple year classes contributing to the fishery with good density of 20-23 inch fish. surface baits at night during spring, or troll deep diving crank baits or spinner rigs

ocation/Species	Forecast	Comments
IARION COUNTY LA	KE 32 acres	(Marion County)
Summertime (mid-J	lune through Sept	ember) oxygen levels are too low to support fish below 11 feet.
Bluegill	Good/Excellent	Good numbers of 8-inch plus fish.
Channel catfish	Good	All sizes available.
Largemouth bass	Fair	High numbers of 11-13 inch fish.
Redear sunfish	Good/Excellent	10-inch plus fish available. Larger fish up shallow in early/mid April for spawn then move out to thermocline depth.
White crappie	Fair/Good	Large fish available; fish laydown trees in the upper end of the lake in spring and brushpiles in late winter and summer.
IARTINS FORK LAK	E 334 acres	(Harlan County)
Bluegill	Good	Good numbers. Most harvested fish are 7.25 inches or less. Fish shallow mud flats in spring or summer.
Channel catfish	Good	Most fish are less than 20 inches; fingerlings were stocked in 2014. Fish at night for better results as water is very clear. There are several good bank fishing areas. Cranks Creek and Martins Fork Creek where first entering the lake when murky can have good numbers of fish.
Crappie	Fair	White crappie are present. Harvested fish are typically 9 to 10 inches. Occasional large fish up to 12-13 inches. Crappie are scattered in shallow water in spring and congregate in tight groups in standing timber in summer.
Largemouth bass	Fair	Numbers over 15 inches are above average at present for the lake. This lake contains 4 species of black bass that may be caught by anglers (largemouth, smallmouth, spotted, and coosa bass). Largemouth bass numbers account for approximately 80% of the black bass in the lake. Fishing success is generally greater at night during most of spring and summer.
Walleye	Fair	This fishery is currently rebuilding with first stocking of native strain walleye in 2013. Stockings have continued in 2014-2015. Numbers of legal-size fish are increasing and look for angler success to steadily improve in coming years.
IAUZY LAKE 81 a	cres (Union Co	ounty, Higginson-Henry WMA)
Bluegill	Fair/Good	Large number of 3-5 inch bluegill, but number of 6-8 inch fish is increasing.

WAULT LAKE 81 a	cres (Union i	County, rigginson-rienty wwa)
Bluegill	Fair/Good	Large number of 3-5 inch bluegill, but number of 6-8 inch fish is increasing.
Channel catfish	Good	Stocked every other year; many 15-22 inch fish in the population.
Largemouth bass	Good	Number of 12-16 inch fish have declined over the last couple of years. Still many 18-21 inch bass. A good number of smaller fish in the population should bring up the number of mid-sized fish as they grow into this size range in the next few years. Fish plastic baits and spinner baits around hardwood brush piles, especially in spring/early summer.
Redear sunfish	Fair/Good	Redear sunfish population continues to improve. Number of fish over 8 inches is increasing.

Location/Species	Forecast	Comments
MCNEELY LAKE	51 acres (Jefferson	n County)
Bluegill	Good	Good numbers of fish through 8 inches.
Channel catfish	Fair	Catchable-size stocked each year; large fish possible.
Common carp	Fair	A few large fish present.
Crappie	Fair	Quality-size fish possible; fish near weed beds and fallen trees, particularly in spring.
Largemouth bas	s Good	Good numbers of fish over the 12-inch size limit. Best cover is water willow, fallen trees and stumps.
Redear sunfish	Good/Excellent	Good numbers of fish through 9 inches; ≥10-inch fish are present.
Warmouth	Fair	Large fish present around weed beds.

И	ETCALFE COUNTY L	<b>_AKE</b> 22 ac	res (Metcalfe County)
	Summertime (mid-Ju	ine through Septe	ember) oxygen levels are too low to support fish below 7 feet.
	Bluegill	Good	High numbers overall with good numbers of 6-inch fish; however, few exceed 8 inches.
	Blue catfish	Fair	Fair numbers overall, but some trophy-size fish (30-inch plus) available. Fish with live or cut bait near creek channel. Stocked every 3rd year.
	Channel catfish	Fair/Good	Fair numbers available. Stocked every other year.
	Crappie	Fair	Smaller-sized (8-9 inch) on average; better sizes available but not overly abundant.
	Largemouth bass	Good/Excellent	Not a high density population; however, trophy fish (20-inch plus) potential consistently above average. Try trout colored jerkbaits in May.
	Rainbow trout	Fair	600 fish stocked in March for a put and take fishery. Water too warm by mid-May to support trout through the summer.

Ī	MILL CREE	K LAKE	109 acres	(Monroe County)

Summertime (mid-June through September) oxygen levels are too low to support fish below 12 feet.

Blue catfish	Fair	Stocked every 3rd year.
Bluegill	Good	Good numbers of 6-inch plus fish.
Channel catfish	Fair	Stocked every other year.
Crappie	Fair	Fair number of 8-9 inch fish. Larger crappie present but at low numbers.
Largemouth bass	Good/Excellent	Good numbers of larger fish (15-inch plus) overall; 18-inch plus fish numbers well above average.
Redear sunfish	Fair	Lower numbers overall, but large fish (10-inch plus) available.
Spotted bass	Poor/Fair	Fair numbers, but slow growing with few fish exceeding 12 inches.

Location/Species	Forecast	Comments
WILL CREEK LAKE		well/Wolfe County – Natural Bridge State Park)
Bluegill	Good	Better numbers recently; excellent numbers of fish over 8 inches. Water is extremely clear and can be difficult to fish.
Channel catfish	Good	Stocked every other year. Note: limb lines (set lines), jug lines, and trot lines are illegal to use at this lake.
Largemouth bass	Good	Similar numbers to previous years. Holds nice 20 inch plus fish, but this lake can be difficult to fish. The timber areas in the upper reaches are the best bet. <b>Note: 15 inch minimum size limit.</b>
Rainbow trout	Excellent	Stocked annually. Best time to fish is within the first few days after stocking.
NOLIN RIVER LAKE	5,795 acres	(Grayson, Edmonson and Hart counties)
Catfish	Good	Channel catfish - good number of 15-20-inch fish and many larger. Fish upper lake (Sportsman's Paradise and Cane Run) and river area in early spring and rocky main lake banks mid-May through June. Flathead catfish - lower numbers than channels but several large fish (30+ lbs) harvested annually. Fish late May through June along the steeper rocky banks.
Crappie	Good	Crappie angling should continue to be good at Nolin in 2016. Good number of 10-11 inch fish in the population. During summer fish deeper brush piles (16-18 feet deep) mid lake for larger fish.
Largemouth bass	Good	Decent number of 15-20 inch largemouth in the population. Anglers should continue to see a good number of quality bass in 2016.
Walleye	Fair	Most fish 14-15 inches with a few in the 20-inch range. In early spring, fish Bacon Creek and upper river with lead head jigs and plastic grubs. In summer, troll Sportsman's Paradise, Conoloway Creek, and the longer main lake points in the Red Rock and Rock Creek areas.
White bass	Excellent	Large number of 11-14-inch white bass in Nolin. Fish Bacon Creek and upper river during spring spawning run with lead head jigs and plastic grubs or live bait. Fish longer main lake points or jumps in summer. Also in summer fish at night under lights near main channel with jigging spoons or live minnows.
OHIO RIVER		
Blue catfish	Good	Good populations in tailwaters and deep holes around outside bends.
Channel catfish	Excellent	Look for silt-free, rocky habitat and embayments.
Common carp	Fair	Large fish possible. Tributaries and backwaters away from current are best areas.
Crappie	Good/Excellent	Backwater areas and creek mouths with brushy habitat are best producers.
Flathead catfish	Good	Potential for very large fish; look for cover such as sunken logs, boulders or other large objects.
Hybrid striped bass	Excellent	Fish up to 6 pounds in and around tailwaters.
Largemouth bass	Fair/Good	Tributaries and backwaters away from current are best areas.
Sauger	Good	Spring and fall below dams with minnows or jigs on bottom.

Location/Species	Forecast	Comments
OHIO RIVER (continu	ıed)	
Smallmouth bass	Fair	Low population in mid/upper river in sections of rocky habitat and tailwaters.
Spotted bass	Fair	Fish the main channel around cover.
Striped bass	Fair	Mid-summer to late fall good for 6-10 pound fish.
Walleye	Fair	Occasionally found in tailwaters.
White bass	Good	Good population.

#### PAINTSVILLE LAKE and TAILWATER 1,139 acres (Morgan and Johnson counties)

Hydrilla is present at Paintsville Lake. Boaters are asked to take measures to prevent the spread of this damaging non-native invasive species by properly inspecting and disinfecting boats, trailers and equipment.

Channel catfish	Fair	Large catfish present in low densities in upper lake. Down-lake large fish are often located deep; fish that are shallow can be found around rock riprap shoreline areas.
Crappie	Good	Most of successful fishing is done on the upper end of the lake into Open Fork and Little Paint Creeks; fish sampled up to 15 inches. Late March can often provide good catches of keeper fish in shallow water near Patoker boat ramp. Look for sunny days at this time of year for best fishing. Lower lake will have larger fish often holding at edges of hydrilla beds.
Largemouth bass	Fair	High survival of young fish due in part to presence of hydrilla which offers ample cover. Stable population with high numbers of fish <12 inches; fish sampled up to 24 inches; slow growth rates; number of fish over 15 inches are increasing and tournament weights are improving. March is popular for catches of lunker fish in standing timber.
Rainbow trout	Fair	Good fishing available in shallow water during winter-early spring after February stocking; some fish present in summer at depth of 20-25 feet. The catch and release portion of the tailwater can offer good quality rainbow and brown trout fishing.
Smallmouth bass	Poor	Poor recruitment; fish sampled up to 22 inches (lower lake areas seem best for larger fish); changes were made in water releases at dam in 2006 to improve summer cool water habitat; spawning benches being placed in lake to aid spawning. Supplemental stockings of native strain fish occurred from 2009-2012.
Spotted bass	Poor	Numbers are down; some large spotted bass present up to 14 inches; best fishing in lower half of lake along main lake shoreline.
Walleye	Fair	Fair numbers through 28 inches; potential for trophy size; look for active fish to be suspended in March; also May and October are good months to fish with live night crawlers on a light jighead fished off the bottom. During summer, drop minnows or nightcrawlers down weedbed edges. Last several year classes are doing well and will provide quite a few keeper fish for 2015.

Location/Species	Forecast	Comments
PENNYRILE LAKE	47 acres (Chi	ristian County)
Bluegill	Good	Some nice bluegill are present as a result of the stunted bass population. Recent surveys found historically high numbers of bluegill > 8 inches. Numerous fish less than 6 inches also present.
Channel catfish	Good	Good numbers of catfish 12-15 inches, and a few larger individuals present as well. Fishing around fallen trees and beaver huts with night crawlers or cheese bait should provide success.
Crappie	Poor	Low overall numbers of crappie, but fish are good size.
Largemouth bass	Fair	This lake has a high number of bass 12 inches and smaller. It is a great place to take kids fishing! Bass are protected between 12-15 inches. Harvesting some smaller ones is allowed, and will help the population.
Redear sunfish	Fair	Numerous fish in the 8-9 inch range. Few redear present greater than 10 inches. Try fishing a small piece of night crawler under a cork near the bottom for best success.
ROUGH RIVER LAKE	and Tailwater*	5,100 acres (Breckinridge and Grayson counties)
Catfish	Good	Good number of 20-25 inch channel catfish with many larger. Lower number of flathead catfish, but trophy fish available. Fish upper lake areas in early spring and steeper rocky banks during mid May-June for spawning fish.
Crappie	Good	Good number of 9-11 inch fish in the population; many 12-13 inch fish as well. Deeper brush piles in mid-lake hold larger fish during summer.
Hybrid striped bass	Excellent	Excellent population of hybrid striped bass. Good number over 20 inches available. Fish upper lake areas in spring. Troll main lake points in summer or fish jumps in main lake (Dam to Tules Creek on north fork and Dam to Little Clifty Creek on south fork). Fish main lake points and sand/mud flats in fall and winter.
Largemouth bass	Good	The number of 15-20-inch bass is higher than the last several years. Bass anglers at Rough should continue to enjoy a good number of quality bass for the next several years.
Tailwater*	Closed	Access to the tailwater will be closed for 3-5 years beginning April 2015 due to repair work being performed on the dam.
SHANTY HOLLOW LA	KE 135 acres	(Warren and Edmonson counties)
Summertime (mid-Ju	une through Sept	tember) oxygen levels are too low to support fish below 10 feet.
Bluegill	Fair	Numbers of 6-inch plus fish up from last couple of years. Fish 8-10 feet on brushpiles in summer for larger fish.
Channel catfish	Fair/Good	All sizes available. Stocked every other year.
Largemouth bass	Fair	Good numbers of 12-14 inch fish.
Redear sunfish	Fair	Fair numbers of 8-inch plus fish. Larger fish up shallow early in the spring then out to thermocline depths by early summer.
White crappie	Fair/Good	Large fish (10-14 inches) available. Fish stakebeds and shallow brushpiles in spring; deeper brush in late winter and summer

deeper brush in late winter and summer.

Location/Species	Forecast	Comments
SMOKY VALLEY LAK	E 36 acres	(Carter County – Carter Caves State Park)
Bluegill	Fair	Slightly better than last season; most less than 7 inches.
Channel catfish	Good	Stocked every other year. Note: limb lines (set lines), jug lines, and trot lines are illegal to use at this lake.
Largemouth bass	Fair	This population of fish is continuing to improve with the more restrictive regulation. There are plenty of chances to catch a good number of fish, but the size is still not there. Note: 12-15 inch protective slot limit; 6 fish daily creel limit, only one over 15 inches.
SPURLINGTON LAKE	36 acres (	Taylor County)
		ember) oxygen levels are too low to support fish below 7 feet.
Bluegill	Good	Good numbers of 6 inch plus fish, with 8-inch plus available.
Channel catfish	Good	All sizes present. Stocked every other year.
Largemouth bass	Excellent	Better than average numbers of larger fish (15-inch plus). Big bass (20-inch plus) opportunities above average.
Redear sunfish	Fair	Lower numbers of fish overall; however, 10-inch plus available.
TAYLORSVILLE LAKE	3,050 acres	(Spencer, Anderson and Nelson counties)
Bluegill	Fair/Good	Good number of fish 6 inches and longer around standing timber in coves.
Catfish	Good/Excellent	Blue catfish population stable; stocked annually since 2002. Fish with live bait, cut bait or crank baits (open water areas). Fish over 30 inches possible. Many channel catfish of all sizes; look for bigger fish in the timber in spring, head of creeks in fall, and points next to channel any time. Numbers of flathead catfish are increasing.
Crappie	Good	Good number of fish at and above the 9-inch size limit. Black crappie now dominate population. Good spawns of white and black crappie were found in 2015. <b>Note: daily creel limit 15 fish.</b>
Hybrid striped bass	Fair/Good	Hybrid striped bass numbers increasing, including numbers of fish greater than 15 inches. In spring and fall, there is good fishing in headwaters. Look for jumps early and late in day July-September.
Largemouth bass	Good	Very good numbers of 12 to 15-inch fish; Good number of 15 to 18-inch fish; best in timber coves; try embayments in spring and fall.
Saugeye	Fair	First year of stocking; harvestable-sized saugeye may be present by fall of 2016.
White bass	Fair/Good	Good numbers in Salt River above the lake in spring (with good flow from rains) and fall; many 10-12 inch fish present.

Location/Species	Forecast	Comments
THREE SPRINGS LAN	KE 30 acres	(Warren County)
Lake is very shallow		
Bluegill	Fair	Fair numbers of 6-8 inch fish. Receives hybrid bluegill stocking in September.
Channel catfish	Good	Good numbers and good size fish (15-inch plus) available. Stocked periodically through the spring and summer.
Largemouth bass	Fair	Numerous fish available; few fish larger than 12 inches. Fish trout colored swim baits after stocking dates and in early to mid-May for larger bass.
Rainbow Trout	Good	Stocked periodically through winter and early spring months (November – March) for put and take fishing. Lake too warm for trout by early-mid May.
Redear	Fair	Fair number of 9-10 inch fish available.
WASHBURN LAKE	18 acres (O	hio County)
Bluegill	Fair	Low number; most fish 6-7 inches.
Channel catfish	Good	Stocked every other year. Good number of fish in the population; many 15-20 inches.
Largemouth bass	Poor	Population dominated by slower growing sub-legal fish. Few fish greater than 12 inches.
Redear sunfish	Fair	Low number; most fish in the 7-8 inch range, but some 9-inch fish available.
WEST FORK DRAKES	S RESERVOIR	88 acres (Simpson County)
Summertime (mid-J	lune through Sep	otember) oxygen levels are too low to support fish below 8 feet.
Bluegill	Good	Good numbers of 6-inch plus fish. Few 8-inch plus fish.
Channel catfish	Good	Numerous sizes available. Fishing best at night near creek channel.
Largemouth bass	Fair/good	Better than average numbers of 12-inch plus fish but few fish larger than 15 inches.
Muskie	Fair/Good	Fishing best below the dam year-round using large inline spinners and jerk baits. Fishing best in lower half of the lake near vegetation beds and brush in winter to early spring.
Redear sunfish	Fair/Good	Good numbers of 6-inch plus fish; few larger than 9 inches.
WOOD CREEK LAKE	672 acres	(Laurel County)
Bluegill	Fair	Moderate numbers of 6-7 inch bluegill.
Diaogiii	ı un	moderate numbers of a ration bluegill.
Channel catfish	Good	Stocked every other year. Consistently produces 12-18 inch fish.
Crappie	Fair	Good density of smaller fish with some fish exceeding 9 inches.
Largemouth bass	Good	Declining big fish population but still has good size range with some fish exceeding 20 inches.
Rainbow trout	Fair	Stocked annually in October and February.

Location/Species	Forecast	Comments
WOOD CREEK LAKE	(continued)	
Redear sunfish	Fair	Moderate density of 8-inch and larger redear are available. Mid and lower lake areas are best.
Spotted bass	Fair	Fair density but most are less than 12 inches. Mid and lower lake areas are best.
YATESVILLE LAKE	2,314 acres	(Lawrence County)
Bluegill	Excellent	Excellent numbers and size distribution throughout the lake; numerous fish 7.0-7.5 inches.
Channel catfish	Excellent	Quality size distribution and densities of catfish exist in all areas of the lake; recent stockings have greatly increased numbers. During fall to early winter many fish are shallow in upper lake; popular lake for jugging and limb lines.
Crappie	Good	Good numbers of 8-15 inch fish. Larger crappie will aggressively go after shad or minnows. Traditional brushy areas and deadfall trees are good for spring spawning fish, with shallow mudflats being good in the fall. This fishery consists only of white crappie.
Flathead catfish	Fair	Good size distribution with fair numbers; rocky shoreline and brush will have better numbers. Good recruitment of young fish in recent years.
Largemouth bass	Good	Good size distribution and good numbers of 15-22 inch fish. Supplemental stockings of fingerling bass were done in 2009, 2012, and 2013 to aid average recruitment. Popular lake for tournament fishing as most of the year produces good results. To avoid boat access congestion problems at ramp, utilize KDFWR website for viewing and scheduling tournaments. Check out KDFWR web page for GPS coordinates of submerged fish habitat structures.
Redear sunfish	Fair	Population is increasing; annual stockings of fish from 2010-2013; larger fish up to 11 inches. Target areas with weedbeds or submerged roadbeds with some rock on them.

#### Fishing Tips for Kentucky and Barkley Lakes

Largemouth bass

During late winter/early spring, look for suspended schools of fish near primary and secondary points. Once located, try using suspended jerkbaits or Alabama rigs. Vary your twitch speeds with your jerkbait until you've found what the fish seem to prefer for the day. Fishing slower is often better, as fish can be slow to take a lure in cold water.

In spring, fish shallow with a Rapala #11 or #13 gold with black back, and floating or shallow diving stick baits. Also try black/brown or black/blue ½-ounce jigs with a #11 or #1 pork frog in the yellow mustard flowers in the backs of embayments. Around woody structure use a spinner bait, or plastic lizards in pumpkin or june bug color.

During summer, fish ledges near river channels with Carolina-rigged and Texas-rigged soft plastics, and deep-diving crank baits. Also fish stumps on the main lake in the mornings and evenings with topwater baits like the zara spook (current is the key to these main lake stumps being productive). Using a 10 inch plastic worm fished along the ledges during the summer months is also good. A good color is plum.

During fall, fish along main lake ledges around stumps and brush with shad or blue/chartreuse color crank baits. Use top water baits (zara spook or devils horse) in the flats of embayments on cloudy days or at dawn and dusk. Use white spinnerbaits around main lake stumps.

The recent dry summers have really brought on the aquatic vegetation. The beds of weeds are extremely productive during the late spring, summer and early fall. Top water baits such as spooks, stick baits, plastic frogs, and buzz baits are great choices. Plastics, such as worms, and spinner baits are also good when fishing the edges of the weed beds.

In spring, fish shallow and over flats with jigs and minnows. Cork fish around buttonball bushes. During summer, white bass anglers trolling small crankbaits near the river channel and primary points in the lake catch many crappie. During fall, fish jigs and minnows next to deep ledges formed by the river channel. Fish may also be caught suspended on flats adjacent to the secondary channel around 10 feet deep near structure. Fall is also a good time to catch black crappie along gravely

shorelines, fishing with a curly tail jig.

During the summer, fish along the main river channel with minnows, cut shad/skipjack herring, and stink baits. An alternative bait option is pieces of Silver carp. One might jump in your boat, or ask a bow fisherman for one.

In spring, fish with leeches or night crawlers around rip rap. During summer, fish with cut shad, chicken livers, minnows, leeches, and worms along the main river channel.

Troll mainlake bars during the late summer and early fall with wiggle-warts or crankbaits, which bump bottom. During fall and winter, use jigs and minnows along the old river channel, along primary points near the river channel, and where a secondary channel meets the old river channel. A good site for fishing is along the river channel just north of the Paris Landing Bridge (US Highway 79).

Fish the jumps from July – September with slab spoons or white roostertails.

Crappie

Blue catfish

Channel catfish

Sauger

White bass

#### Fishing Tips for Kentucky and Barkley Tailwaters

Blue and channel catfish The best time is in May-June using cut bait or live bait just off the bottom.

Sauger Fish ¾-oz or 1-oz jigs with minnows bumping bottom in back eddies and

near current. Trolling is also good during the winter and early spring.

Best fishing is between December and February.

Striped bass and hybrids The best time is from April-June using live bait (skipjack herring) or 4-

inch sassy shad baits with a 1-2 ounce head in pearl or chartreuse color. The amount of weight depends on the current and should allow baits to bump the bottom. Live bait can be caught in the tailwaters using a cast

net or a skipjack rod.

# Fishing Forecast Cheat Sheet (Summary of lakes forecasting good or excellent fishing for each species listed)

	Largemouth	Smallmouth	Spotted		Redear		Rainbow	Brown	
Water body	bass	bass	bass	Bluegill	sunfish	Crappie	trout	trout	Muskellunge
Barkley Lake/tailwater	<b></b>			<b>_</b>	_	<b></b>			
Barren River									<b></b>
Barren River Lake									
Beaver Lake				-		•			
Benjy Kinman Lake	•			•					
Bert Combs Lake	./			./			1		
Beshear Lake	./			·			*		
Beulah Lake							-/		+
Boltz Lake	V						·		-
Briggs Lake	V								
					V				
Buckhorn Lake/tailwater									+
Bullock Pen Lake									
Cannon Creek Lake									
Carpenter Lake									
Carr Creek Lake	<b>√</b>		<b>V</b>			<b></b>			
Cave Run Lake	<b>✓</b> ,		$\checkmark$						$\checkmark$
Cedar Creek Lake	$\checkmark$			<b>✓</b>	<b>√</b>	$\checkmark$			
Clear Creek Lake				$\sqrt{}$	$\checkmark$				
Corinth Lake									
Cumberland Lake	<b>√</b>		<b>√</b>	<b></b>		<b>√</b>			
Cumberland Tailwater		·	Ť	Ť		Ť	<b>√</b>	<b>√</b>	
Dale Hollow Lake			<b></b>						
Dewey Lake	1	¥	*			1			
Elkhorn Creek	./	-/							
Elmer Davis Lake		V			./				
Fagan Branch Lake	V								
Fishtrap Lake/tailwater					V				1
		V							
Grayson Lake	V								
Green River		<b>✓</b>							V .
Green River Lake	V_		<b>✓</b>			<b>✓</b>			<b>✓</b>
Greenbo Lake							<b>√</b>		
Guist Creek Lake	<b>√</b>								
Herrington Lake				<b></b>					
Kentucky Lake/tailwater	$\checkmark$	<b>√</b>		<b>_</b>	$\checkmark$	<b></b>			
Kentucky River									<b>V</b>
Kincaid Lake	<b>√</b>								
Lake Malone									
Lake Reba				_/					
Lake Wilgreen	./			-	./				
Laurel River Lake			./	-/-	· ·				
Marion County Lake	V			<u> </u>					
					V				
Martin's Fork Lake				<b>├</b> ✓					+
Mauzy Lake	<b>V</b>								+
McNeely Lake	<b>V</b>			V_	✓				1
Metcalfe County Lake	<b>√</b> ,			<b>✓</b> ,					
Mill Creek Lake (Monroe Co.)	<b>V</b> ,			<u></u>					1
Mill Creek Lake (Powell/Wolfe Co.)				$\checkmark$			$\checkmark$		
Nolin River Lake						$\overline{}$			
Ohio River						$\overline{}$			
Paintsville Lake/tailwater						<b></b>			
Pennyrile Lake				<b></b>					
Rough River Lake						<b></b>			
Smoky Valley Lake	•								1
Spurlington Lake	./			./					1
Taylorsville Lake	\ \ \ \ \ \ \ \ \ \ \ \ \ \ \ \ \ \ \			<del>- v</del>				-	+
Three Springs Lake	V			-	-			-	+
Washburn Lake	<del>  </del>			<del>                                     </del>			V	<b> </b>	+
				_	-			-	+
West Fork Drakes Reservoir				<b>✓</b>					+
Wood Creek Lake									
Yatesville Lake	$\checkmark$	nal species		$\checkmark$		$\overline{}$			

<sup>\*\*</sup> Additional species can be found on following page.

# Fishing Forecast Cheat Sheet (Summary of lakes forecasting good or excellent fishing for each species listed)

	Striped	Hybrid striped	White			Channel	Blue	Flathead
Water body	bass	bass	bass	Walleye	Sauger	catfish	catfish	catfish
Barkley Lake/tailwater				1			•/	
Barren River		/				./		
Barren River Lake	1	1				·/	-/	./
Beaver Lake	1	· ·				./	<u>v</u>	_ v
Benjy Kinman Lake	1					Y		
Bert Combs Lake	1			-				
Beshear Lake	+							
Beulah Lake	1					V		
Boltz Lake	1							
Briggs Lake	1							
	┥ ├───							
Buckhorn Lake/tailwater Bullock Pen Lake	┥┝───			-				V
	┥┝──			-				
Cannon Creek Lake	┥┝──			-				
Carpenter Lake	4					<b>✓</b>		
Carr Creek Lake	4							
Cave Run Lake	4					<b>√</b>		
Cedar Creek Lake								
Clear Creek Lake	<b>↓</b>							ļ
Corinth Lake				<u> </u>		V.		
Cumberland Lake	$\overline{}$					$\overline{}$		
Cumberland Tailwater	] [		$\overline{}$					
Dale Hollow Lake	1	l i	*					
Dewey Lake							$\overline{}$	<b>V</b>
Elkhorn Creek							*	
Elmer Davis Lake	1					-		
Fagan Branch Lake	1					./		
Fishtrap Lake/tailwater	+					V_		
Grayson Lake	+	- V					<u>v</u>	<del>- v</del>
Green River	<del>                                     </del>	V .						- V
Green River Lake	<del>                                   </del>	V						
	┥┝───			-				
Greenbo Lake	┥							
Guist Creek Lake	┥					<b>✓</b>		
Herrington Lake	<del>                                     </del>							
Kentucky Lake/tailwater					,	<b>√</b>		
Kentucky River	4							
Kincaid Lake								
Lake Malone						<b></b>		
Lake Reba						<b>√</b>		
Lake Wilgreen								
Laurel River Lake				<b>V</b>				
Marion County Lake						<b>_</b>		
Martin's Fork Lake								
Mauzy Lake								
McNeely Lake	1							
Metcalfe County Lake	1	1						
Mill Creek Lake (Monroe Co.)	1	1		<b>†</b>				
Mill Creek Lake (Powell/Wolfe Co.)	1	1		1		./		
Nolin River Lake	1	1	./	1		./		<b></b>
Ohio River	1 🗕 —	<del>                                     </del>	<u>v</u>	+	./	./		- ×
Paintsville Lake/tailwater	1	<del></del>		+		V		<del>                                     </del>
Pennyrile Lake	┨ ├───	+		1				1
	┨├──	<del>                                     </del>		-		- V		-
Rough River Lake	┪┝───	<b>✓</b>		+		- V		<b> </b>
Smoky Valley Lake	┨┝───	1		1		<b>V</b>		<del>                                     </del>
Spurlington Lake	<b>↓</b>					<b>V</b> ,		<u> </u>
Taylorsville Lake	<b>↓</b>			1		<b>V</b> ,		
Three Springs Lake		1		1	I	<b>✓</b>		
Washburn Lake						<b>V</b>		
West Fork Drakes Reservoir						<b>-</b>		
						\ \ \ \		